



5.07-101  
10.07-1020 I

304988

"CEPHALOSPORIN DERIVATIVES, THEIR PREPARATION AND  
COMPOSITIONS CONTAINING THEM"

The present invention relates to a series of new cephalosporin compounds which are particularly suitable for oral administration, to processes and intermediates for preparing these compounds and to compositions containing the compounds.

Although many cephalosporin derivatives which exhibit excellent antibacterial activity have been discovered, most of them are for parenteral administration. However, except where massive doses of an antibiotic are to be administered quickly, the preferred route of administration is oral, as oral preparations can be administered by the patient himself without the need for trained supervision or assistance. Unfortunately, of the many cephalosporin derivatives discovered, very few possess a combination of superior antibacterial activity, broad antibacterial spectrum against both gram-positive and gram-negative bacteria (especially against Staphylococcus aureus) and the ability to be absorbed efficiently through the digestive tract.

For example, cephalothin, cefazolin and cefmetazole are widely used for parenteral administration, particularly by injection. However, when these compounds

09/25/81	304988	1 101	65.00CK
09/25/81	304988	1 102	106.00CK

2

are administered orally, only about 5% of the dose administered is recovered in the urine, showing their poor absorption through the digestive tract and their unsuitability for oral administration. This is thought  
5 to be due to the strong dissociation of the carboxy group at the 4- position (i.e. the low pKa value) and the strong acidity.

Because of this, many efforts have been made to improve the absorption of cephalosporin derivatives  
10 through the digestive tract by esterifying the 4- carboxy group but almost all such efforts have failed to obtain cephalosporin derivatives which are well absorbed through the digestive tract and which are therefore useful for oral administration; as described hereafter, in the one  
15 instance where absorption through the digestive tract has been significantly improved, the resulting compound lacks the desired broad antibacterial spectrum.

For example, the Journal of Antibiotics, 32 No. 11, 1155 (1979) discloses that the absorption of  
20 cefamandol through the digestive tract is not improved by esterification to prepare the acetoxymethyl ester, since this ester is only sparingly soluble in water. Although absorption of the ester through the digestive tract can be improved to a limited extent by administration  
25 of the ester in solution in certain organic solvents (such as propylene glycol), this is not a particularly good


3

solution to the problem.

The Journal of Medicinal Chemistry, 22, 657  
(1979), on the other hand, reports that the absorption  
through the digestive tract of another ester of a  
5 cephalosporin which is readily soluble in water, is not  
significantly improved due to chemical instability of  
the ester.

Furthermore, it is known that, in general, lower  
alkyl and benzhydryl esters of cephalosporins possess, in  
10 themselves, almost no antibacterial activity and that  
they are not hydrolyzed in vivo (which might otherwise  
convert them to an active acid) and hence they are not  
of value for therapeutic use, although they may be useful  
as synthetic intermediates.

15 Of the various cephalosporin derivatives known,  
one known class has a 2-(2-aminothiazol-4-yl)-2-alkoxy-  
iminoacetamido group at the 7- position and may be  
represented by the following formula:



(in which B, D and E are substituents).

For example, Japanese Patent Application  
Kokai (i.e. as laid-open to public inspection) No.  
*which corresponds to USP 4,098,888*  
149296/76, discloses the following compounds:

- 5 (a) 7-[2-(2-aminothiazol-4-yl)-2-methoxyimino-  
acetamido]-3-methyl-3-cephem-4-carboxylic acid;
- (b) 3-acetoxymethyl-7-[2-(2-aminothiazol-4-yl)-  
2-methoxyiminoacetamido]-3-cephem-4-carboxylic acid;  
and
- 10 (c) 7-[2-(2-aminothiazol-4-yl)-2-methoxyimino-  
acetamido]-3-[(1-methyl-1H-tetrazol-5-yl)thiomethyl]-3-  
cephem-4-carboxylic acid.

We have discovered that the percentage recovery of these  
compounds in urine (which is a measure of their suit-  
15 ability for oral administration) is only 3.2%, 1.5% and  
2%, for compounds (a), (b) and (c), respectively, these  
compounds are, accordingly, unsuitable for oral  
administration.

Likewise, Japanese Patent Application Kokai  
20 No. 86188/81 ~~(which was published after the priority date~~

*a* ~~of the present Application)~~ discloses 7-[2-(2-amino-thiazol-4-yl)-2-methoxyiminoacetamido]-3-methoxymethyl-3-cephem-4-carboxylic acid (d), which is the free carboxylic acid corresponding to certain of the compounds  
 5 of the present invention. We have, however, found that the recovery rate in urine of compound (d) is only 5.5% and it is, therefore, unsuitable for oral administration. The Specification also discloses certain esters, particularly the t-butyl and benzhydryl esters, of  
 10 cephalosporin compounds related to compound (d). However, as stated above, such esters are not believed to be readily convertible in vivo to the corresponding carboxylic acid and, as a result, may not be effective in actual use.

15

*which corresponds to USP 4,278,793*  
 Japanese Patent Applications Kokai No. 9296/79  
*which corresponds to USP 4,278,671*  
 and 34795/78, disclose the following pivaloyloxymethyl esters:

*a* (e) pivaloyloxymethyl 3-acetoxymethyl-7-[2-(2-amino-thiazol-4-yl)-2-methoxyiminoacetamido]-3-cephem-4-  
 20 carboxylate and

(f) pivaloyloxymethyl 7-[2-(2-aminothiazol-4-yl)-2-methoxyiminoacetamido]-3-(1-methyl-1H-tetrazol-5-yl)-thiomethyl-3-cephem-4-carboxylate.

7

13 We have also found that the recovery rate in urine of these compounds is only 8% and 14% for compounds (e) and (f), respectively, and these compounds also are unsuitable for oral administration.

5 P Comparing the recovery rates of compounds (a), (b) and (c) with the recovery rates of compounds (e) and (f), the results are rather surprising, since it is known that the absorption of ampicillin through the digestive tract is considerably improved by converting it to the  
10 pivaloyloxymethyl ester.

The above-mentioned Japanese Patent Application Kokai No. 34795/78 also discloses pivaloyloxymethyl  
29 7-[2-(2-aminothiazol-4-yl)-2-methoxyiminoacetamido]-3-methyl-3-cephem-4-carboxylate, hereinafter referred to  
15 as "compound (g)". We have carried out extensive studies of this compound and have found that it exhibits very good recovery in urine, at a level almost comparable with that of the compounds of the present invention, thus suggesting that it may well be suitable for oral  
20 administration. However, as will be shown hereafter, compound (g), when administered orally, is hydrolyzed and converted in vivo to the corresponding carboxylic acid which, in turn, has poor activity against Staphylococcus aureus. Failure to inhibit the growth  
25 of this bacterium, which is perhaps one of the most

8

important from the clinical point of view, could be a disadvantage in actual use.

It is, accordingly, clear from the above discussion that preparation of a cephalosporin derivative which meets the triple requirements of good absorption through the digestive tract, high antibacterial activity and a broad antibacterial spectrum, is not a simple matter. The cephalosporin nucleus includes many points at which different substituents may be introduced and the introduction of a particular substituent to improve one property may adversely affect other properties in a quite unpredictable way. Moreover, it has clearly been demonstrated that, even where a particular chemical modification is known to improve the properties of one particular compound (e.g. as with the preparation of the pivaloyloxymethyl ester to improve the absorption of ampicillin), this is not any indication that a similar modification will similarly improve the properties of any other compound.


We have now surprisingly discovered a limited class of cephalosporin derivatives which can be administered orally as they are readily absorbed through the digestive tract and which are then readily

9

8.

hydrolyzed and converted in vivo to the corresponding carboxylic acid which, in turn, shows quite outstanding activity against both gram- positive and gram- negative bacteria.

5 Accordingly, the present invention consists in compounds of formula (I):



13 in which:

13  $R^1$  represents a methyl group or an ethyl group;

10  $R^2$  represents a hydrogen atom or a methyl group; and

13  $R^3$  represents a  $C_1-C_5$  alkyl or alkoxy group;

13 and pharmaceutically acceptable acid addition salts thereof.

10



The invention also provides a pharmaceutical composition comprising, as active ingredient, one or more of the compounds of the invention in admixture with a pharmaceutically acceptable carrier or diluent.

5           The invention also provides a variety of processes for preparing the compounds of the invention.

10           In the compounds of formula (II), when  $R^3$  represents an alkyl group having from 1 to 5 carbon atoms, it is preferably a methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl, t-butyl, pentyl, isopentyl or t-pentyl group, most preferably a t-butyl group.  $R^3$  most preferably represents an alkyl group having from 1 to 5 carbon atoms when  $R^2$  represents a hydrogen atom.

15           When  $R^3$  represents an alkoxy group having from 1 to 5 carbon atoms, it is preferably a methoxy, ethoxy, propoxy, isopropoxy, butoxy, isobutoxy, sec-  
20           butoxy, t-butoxy, pentyloxy or 1-ethylpropoxy group, most preferably an ethoxy group.  $R^3$  most preferably represents an alkoxy group having from 1 to 5 carbon atoms when  $R^2$  represents a methyl group.

Examples of compounds of the invention are given in the following list; the compounds are hereafter

11

identified by the numbers assigned to them in the list.

1. Acetoxymethyl 7-[2-(2-aminothiazol-4-yl)-2-methoxyiminoacetamido]-3-methoxymethyl-3-cephem-4-carboxylate
2. Propionyloxymethyl 7-[2-(2-aminothiazol-4-yl)-2-methoxyiminoacetamido]-3-methoxymethyl-3-cephem-4-carboxylate
3. 1-Acetoxyethyl 7-[2-(2-aminothiazol-4-yl)-2-methoxyiminoacetamido]-3-methoxymethyl-3-cephem-4-carboxylate
4. Propionyloxymethyl 7-[2-(2-aminothiazol-4-yl)-2-ethoxyiminoacetamido]-3-methoxymethyl-3-cephem-4-carboxylate
5. Isopropionyloxymethyl 7-[2-(2-aminothiazol-4-yl)-2-ethoxyiminoacetamido]-3-methoxymethyl-3-cephem-4-carboxylate
6. Butyryloxymethyl 7-[2-(2-aminothiazol-4-yl)-2-ethoxyiminoacetamido]-3-methoxymethyl-3-cephem-4-carboxylate
7. 1-Propionyloxyethyl 7-[2-(2-aminothiazol-4-yl)-2-ethoxyiminoacetamido]-3-methoxymethyl-3-cephem-4-

carboxylate

P 8. Isobutyryloxymethyl 7-[2-(2-aminothiazol-4-yl)-<sup>8</sup>  
2-methoxyiminoacetamido]-3-methoxymethyl-3-cephem-4-<sup>9</sup>  
carboxylate

5 P 9. Pivaloyloxymethyl 7-[2-(2-aminothiazol-4-yl)-<sup>8</sup>  
2-methoxyiminoacetamido]-3-methoxymethyl-3-cephem-4-<sup>9</sup>  
carboxylate

P 10. Pivaloyloxymethyl 7-[2-(2-aminothiazol-4-yl)-2-<sup>8</sup>  
ethoxyiminoacetamido]-3-methoxymethyl-3-cephem-4-<sup>9</sup>  
10 carboxylate

P 11. Isovaleryloxymethyl 7-[2-(2-aminothiazol-4-yl)-<sup>8</sup>  
2-methoxyiminoacetamido]-3-methoxymethyl-3-cephem-4-<sup>9</sup>  
carboxylate

P 12. 1-Pivaloyloxyethyl 7-[2-(2-aminothiazol-4-yl)-<sup>8</sup>  
15 2-ethoxyiminoacetamido]-3-methoxymethyl-3-cephem-4-<sup>9</sup>  
carboxylate

P 13. Methoxycarbonyloxymethyl 7-[2-(2-aminothiazol-<sup>8</sup>  
4-yl)-2-methoxyiminoacetamido]-3-methoxymethyl-3-cephem-<sup>9</sup>  
4-carboxylate

P<sub>1</sub> 14. 1-Methoxycarbonyloxyethyl 7-[2-(2-aminothiazol-  
4-yl)-2-ethoxyiminoacetamido]-3-methoxymethyl-3-cephem-  
4-carboxylate 9

P<sub>1</sub> 15. Ethoxycarbonyloxymethyl 7-[2-(2-aminothiazol-  
5 4-yl)-2-methoxyiminoacetamido]-3-methoxymethyl-3-  
cephem-4-carboxylate 9

P<sub>1</sub> 16. 1-Ethoxycarbonyloxyethyl 7-[2-(2-aminothiazol-  
4-yl)-2-methoxyiminoacetamido]-3-methoxymethyl-3-cephem-  
4-carboxylate 9

P<sub>1</sub> 17. 1-Ethoxycarbonyloxyethyl 7-[2-(2-aminothiazol-  
4-yl)-2-ethoxyiminoacetamido]-3-methoxymethyl-3-cephem-  
4-carboxylate 9

P<sub>1</sub> 18. Propoxycarbonyloxymethyl 7-[2-(2-aminothiazol-  
4-yl)-2-ethoxyiminoacetamido]-3-methoxymethyl-3-cephem-  
15 4-carboxylate 9

P<sub>1</sub> 19. 1-Isopropoxycarbonyloxyethyl 7-[2-(2-aminothiazol-  
4-yl)-2-methoxyiminoacetamido]-3-methoxymethyl-3-cephem-  
4-carboxylate 9

P<sub>1</sub> 20. 1-Butoxycarbonyloxyethyl 7-[2-(2-aminothiazol-  
20 4-yl)-2-ethoxyiminoacetamido]-3-methoxymethyl-3-cephem-  
4-carboxylate. 9

14

21. Isobutoxycarbonyloxymethyl 7-[2-(2-aminothiazol-4-yl)-2-methoxyiminoacetamido]-3-methoxymethyl-3-cephem-4-carboxylate
22. 1-sec-Butoxycarbonyloxyethyl 7-[2-(2-aminothiazol-4-yl)-2-methoxyiminoacetamido]-3-methoxymethyl-3-cephem-4-carboxylate
23. 1-(1-Ethylpropoxycarbonyloxy)ethyl 7-[2-(2-aminothiazol-4-yl)-2-methoxyiminoacetamido]-3-methoxymethyl-3-cephem-4-carboxylate
24. 1-(1-Ethylpropoxycarbonyloxy)ethyl 7-[2-(2-aminothiazol-4-yl)-2-ethoxyiminoacetamido]-3-methoxymethyl-3-cephem-4-carboxylate
25. 3,3,3-Trimethylpropionyloxymethyl 7-[2-(2-aminothiazol-4-yl)-2-methoxyiminoacetamido]-3-methoxymethyl-3-cephem-4-carboxylate.

Of the compounds listed above, Compounds No. 9, 10, 16 and 17 are most preferred.


As indicated in the formula, the compounds of formula (I) of the present invention are in the syn-form which has been found to have much stronger antibacterial activity than the corresponding anti-isomers.

The compounds of formula (I) will form acid addition salts with various acids and the invention thus also includes such salts with pharmaceutically acceptable acids, for example inorganic acids (such as hydrochloric acid, sulphuric acid or phosphoric acid) or organic acids (such as methanesulphonic acid, benzenesulphonic acid or malonic acid). Of the acid addition salts, the hydrochlorides are most preferred.

The compounds of the present invention may be prepared by a number of methods, for example those described below.

#### Method 1


Compounds of formula (I) can be prepared by reacting a compound of formula (II):



(in which  $\text{R}^4$  represents an amino group or a protected amino group, and  $\text{R}^1$  is as defined above) or a reactive

15.

derivative of said compound of formula (II) with a compound of formula (III):



(in which  $R^2$  and  $R^3$  are as defined above) and, if  
5 necessary, deprotecting the group  $R^4$ .

9  
10 In the above compounds of formula (II), preferred amino-protecting groups included in  $R^4$  are those groups which may readily be removed to restore a free amino group, for example the trityl, formyl, t-butoxycarbonyl or 2-ethoxycarbonyl-1-methylvinyl groups, which may be removed by treatment with an acid, the 2,2,2-trichloroethoxycarbonyl group, which may be removed by reduction, the 2-methanesulphonylethoxycarbonyl group, which may be removed by treatment with an alkali, or the chloroacetyl  
15 group, which may be removed by treatment with thiourea.

The carboxylic acid of formula (II) may itself be used in the free form or it may be used in the form of a reactive derivative. Suitable reactive derivatives include the acid halide, the acid anhydride, mixed acid

anhydrides, reactive esters, reactive amides and the acid azide. Preferred mixed acid anhydrides include mixed acid anhydrides with mono-(lower alkyl)carbonates, such as monomethyl carbonate or monoisobutyl carbonate, and mixed acid anhydrides with lower alkanolic acids, such as pivalic acid or trichloroacetic acid. Preferred reactive esters include the p-nitrophenyl ester, the pentachlorophenyl ester and the N-hydroxyphthalimide ester.

Where the compound of formula (II) is employed in the form of the free acid, we prefer to carry out the reaction in the presence of a condensing agent. Examples of suitable condensing agents include: di-substituted carbodiimides, such as dicyclohexylcarbodiimide; imidazolides, such as carbonyldiimidazole or thionyl-diimidazole; N-ethoxycarbonyl-2-ethoxy-1,2-dihydroquinoline; or a Vilsmeier reagent prepared from dimethylformamide and, for example, phosphorus oxychloride or thionyl chloride.

Where a reactive derivative of the acid (II) is employed, the use of such a condensing agent is not necessary; however, for certain reactive derivatives, it may be desirable to carry out the reaction in the presence of a base. Examples of suitable bases include: alkali metal compounds, such as sodium bicarbonate, potassium



bicarbonate, sodium carbonate or potassium carbonate;  
or aliphatic, aromatic or nitrogen- containing hetero-  
cyclic bases, such as triethylamine, N,N-dimethylaniline,  
N,N-diethylaniline, N-methyldiethylamine, N-methyldiethyl-  
5 idine, pyridine, collidine or lutidine.

The reaction of the acid (II) or its reactive  
derivative with the compound of formula (III) is preferably  
effected in the presence of a solvent, the nature of  
which is not critical, provided that it has no adverse  
10 effect upon the reaction. Preferred solvents include  
inert organic solvents (such as acetone, methyl ethyl  
ketone, tetrahydrofuran, dioxan, ethyl acetate, chloroform,  
methylene chloride, acetonitrile, dimethylformamide  
or dimethylsulphoxide) or a mixture of such a solvent  
15 and water.

There is no particular limitation on the reaction  
temperature, but we normally prefer to conduct the  
reaction at ambient temperature or with cooling. The  
time required for the reaction will vary, depending mainly  
20 upon the method of acylation and the reaction temperature,  
but usually the reaction will require a period which may  
vary from several tens of minutes to several tens of  
hours.

After completion of the reaction, the reaction

19

product may be recovered from the reaction mixture by conventional means. For example, if a water-miscible solvent is employed, the solvent is preferably removed by distillation under reduced pressure and the residue is dissolved in a water-immiscible solvent. The resulting solution is then washed with an acid and a base and dried, after which the solvent is distilled off to give the desired product. If a water-immiscible solvent is employed for the reaction, the reaction mixture is washed with an acid or a base and dried, after which the solvent is distilled off. The product thus obtained may, if necessary, be further purified by conventional means, for example by chromatographic techniques.


The reaction required to remove the protecting group, if  $R^4$  represents a protected imino group, is, as mentioned above, conventional and will vary depending upon the particular protecting group chosen. After removal of the protecting group, the desired product may be recovered from the reaction mixture and purified, e.g. as suggested above, to give the desired compound of formula (I).

CL Method 2:


f Compounds of formula (I) may be obtained by

20

reacting a compound of formula (IV):



1021K (in which  $R^1$  and  $R^4$  are as defined above) or a reactive derivative thereof with a compound of formula (V):



513 (in which X represents a halogen atom, such as a chlorine, bromine or iodine atom, preferably an iodine atom, and  $R^2$  and  $R^3$  are as defined above) and then, if necessary, deprotecting the group represented by  $R^4$ .

10 Preferred reactive derivatives of the compound of formula (IV) are salts, for example salts with a metal (such as sodium or potassium) or with an organic amine (such as triethylamine). Where the free acid (IV) is employed, the reaction is preferably effected in the

21

presence of an acid-binding agent, which may be organic or inorganic, for example potassium carbonate, sodium carbonate, sodium bicarbonate, triethylamine, dicyclohexylamine, pyridine or N,N-dimethylaniline.

5           The reaction is preferably effected in the presence of a solvent, the nature of which is not critical, provided that it has no adverse effect upon the reaction. Suitable solvents include dimethylformamide, dimethylacetamide, dimethyl sulphoxide, hexamethylphosphoric  
10 triamide or acetonitrile; a mixture of two or more such solvent may be employed, as may a mixture of one or more of these solvents with one or more other inert organic solvents. The reaction may be effected over a wide range  
15 at ambient temperature or with cooling. The time required for the reaction may vary from a period of several minutes to several hours.


          After completion of the reaction, the reaction mixture is preferably diluted with a water-immiscible  
20 solvent, washed successively with an aqueous solution of potassium bisulphate and an aqueous basic solution and then dried, after which the solvent is distilled off to give the desired product. This product may be  
25 further purified by conventional means, for example by chromatographic techniques.

22

Where  $R^4$  represents a protected amino group, it may be converted to a free amino group as described in Method 1.

CL Method 3

5  $\rho$  Compounds of formula (I) may be obtained by reacting a compound of formula (VI):



$\beta$  (in which  $R^5$  represents an alkyl group or an aryl group, and  $R^1$ ,  $R^2$  and  $R^3$  are as defined above) with  
 10 thiourea. Compounds of formula (VI) are new and themselves form part of the present invention.

$\rho$  In the compounds of formula (VI), when  $R^5$  represents an alkyl group, it is preferably an alkyl group having from 1 to 6 carbon atoms, for example a  
 15 methyl, ethyl, propyl, isopropyl, butyl, isobutyl, pentyl or hexyl group, more preferably a methyl or ethyl group. When  $R^5$  represents an aryl group, it is preferably a substituted or unsubstituted phenyl or naphthyl

group. In the case of substituted groups, there may be one or more substituents, normally from 1 to 5 substituents, and they may be the same or different. Suitable substituents include  $C_1-C_4$  alkyl groups (e.g. methyl, ethyl, propyl, isopropyl or butyl),  $C_1-C_4$  alkoxy groups (e.g. methoxy, ethoxy, propoxy, isopropoxy, butoxy or isobutoxy) and halogen atoms (e.g. chlorine, bromine or fluorine atoms). The most preferred aryl groups represented by  $R^5$  are the phenyl and p-methyl-phenyl groups.

Representative examples of compounds of formula (VI) include:

26. Acetoxymethyl 7-(2-methoxyimino-3-oxo-4-p-toluenesulphonyloxybutyrylamino)-3-methoxymethyl-3-cephem-4-carboxylate

27. 1-Acetoxyethyl 7-(2-ethoxyimino-3-oxo-4-p-toluenesulphonyloxybutyrylamino)-3-methoxymethyl-3-cephem-4-carboxylate

28. Propionyloxymethyl 7-(4-benzenesulphonyloxy-2-methoxyimino-3-oxobutyrylamino)-3-methoxymethyl-3-cephem-4-carboxylate

29. 1-Propionyloxyethyl 7-(4-methanesulphonyloxy-2-methoxyimino-3-oxobutyrylamino)-3-methoxymethyl-3-cephem-4-carboxylate

24

- P 30. 1-Butyryloxyethyl 7-(4-ethanesulphonyloxy-2-ethoxy-  
imino-3-oxobutyrylamino)-3-methoxymethyl-3-cephem-4-  
carboxylate
- P 31. Isobutyryloxymethyl 7-(2-methoxyimino-3-oxo-  
5 4-p-toluenesulphonyloxybutyrylamino)-3-methoxymethyl-  
3-cephem-4-carboxylate
- P 32. Pivaloyloxymethyl 7-(2-methoxyimino-3-oxo-4-  
p-toluenesulphonyloxybutyrylamino)-3-methoxymethyl-3-  
cephem-4-carboxylate
- 10 P 33. Pivaloyloxymethyl 7-(2-ethoxyimino-3-oxo-4-  
p-toluenesulphonyloxybutyrylamino)-3-methoxymethyl-3-  
cephem-4-carboxylate
- P 34. Pivaloyloxymethyl 7-(4-benzenesulphonyloxy-2-  
methoxyimino-3-oxobutyrylamino)-3-methoxymethyl-3-  
15 cephem-4-carboxylate
- P 35. Pivaloyloxymethyl 7-(4-methanesulphonyloxy-2-  
methoxyimino-3-oxobutyrylamino)-3-methoxymethyl-3-  
cephem-4-carboxylate
- P 36. Pivaloyloxymethyl 7-(4-ethanesulphonyloxy-2-  
20 ethoxyimino-3-oxobutyrylamino)-3-methoxymethyl-3-cephem-  
4-carboxylate

- P<sub>1</sub> 37. 1-Pivaloyloxyethyl 7-(2-methoxyimino-3-oxo-  
4-p-toluenesulphonyloxybutyrylamino)-3-methoxymethyl-  
3-cephem-4-carboxylate
- 5 P<sub>1</sub> 38. Methoxycarbonyloxymethyl 7-(4-methanesulphonyl-  
oxy-2-methoxyimino-3-oxobutyrylamino)-3-methoxymethyl-  
3-cephem-4-carboxylate
- P<sub>1</sub> 39. Ethoxycarbonyloxymethyl 7-(4-benzenesulphonyl-  
oxy-2-ethoxyimino-3-oxobutyrylamino)-3-methoxymethyl-  
3-cephem-4-carboxylate
- 10 P<sub>1</sub> 40. 1-Ethoxycarbonyloxyethyl 7-(2-methoxyimino-3-  
oxo-4-p-toluenesulphonyloxybutyrylamino)-3-methoxy-  
methyl-3-cephem-4-carboxylate
- P<sub>1</sub> 41. 1-Ethoxycarbonyloxyethyl 7-(4-methanesulphonyl-  
oxy-2-methoxyimino-3-oxobutyrylamino)-3-methoxymethyl-  
15 3-cephem-4-carboxylate
- P<sub>1</sub> 42. 1-Ethoxycarbonyloxyethyl 7-(2-ethoxyimino-3-  
oxo-4-p-toluenesulphonyloxybutyrylamino)-3-methoxy-  
methyl-3-cephem-4-carboxylate
- P<sub>1</sub> 43. Isopropoxycarbonyloxymethyl 7-(2-methoxyimino-  
20 3-oxo-4-p-toluenesulphonyloxybutyrylamino)-3-methoxy-




methyl-3-cephem-4-carboxylate.

44. 1-Butoxycarbonyloxyethyl 7-(4-benzenesulphonyl-  
oxy-2-methoxyimino-3-oxobutyrylamino)-3-methoxymethyl-  
3-cephem-4-carboxylate

45. 1-(1-Ethylpropoxycarbonyloxy)ethyl 7-(2-  
methoxyimino-3-oxo-4-p-toluenesulphonyloxybutyryl-  
amino)-3-methoxymethyl-3-cephem-4-carboxylate

46. 3,3,3-Trimethylpropionyloxymethyl 7-(2-  
methoxyimino-3-oxo-4-p-toluenesulphonyloxybutyryl-  
amino)-3-methoxymethyl-3-cephem-4-carboxylate


Compounds of formula (VI) may be prepared,  
for example, by reacting a compound of formula (VII):



702707

(in which  $\text{R}^1$  and  $\text{R}^5$  are as defined above) or a reactive  
derivative thereof with a compound of formula (III):



27



PS (in which  $R^2$  and  $R^3$  are as defined above). Compounds of formula (VII) are new and also part of the present invention. Representative examples of compounds of formula (VII) include:

- 5 *P* 47. 2-Methoxyimino-3-oxo-4-*p*-toluenesulphonyloxy-butyrlic acid
- P* 48. 2-Ethoxyimino-3-oxo-4-*p*-toluenesulphonyloxy-butyrlic acid
- 10 *P* 49. 4-Benzenesulphonyloxy-2-methoxyimino-3-oxo-*C* butyrlic acid
- P* 50. 4-Methanesulphonyloxy-2-methoxyimino-3-oxo-*C* butyrlic acid
- P* 51. 4-Ethanesulphonyloxy-2-ethoxyimino-3-oxo-*C* butyrlic acid.

Compounds of formula (VII) may, for example, be prepared by the series of reactions illustrated in the following reaction scheme for the preparation of the compound in which  $R^5$  represents a *p*-tolyl group and  $R^3$  represents a methyl group:



70290f

29

*P* In the reaction to prepare the compound of formula (VI), the free acid of formula (VII) may be used as such or a reactive derivative of this free acid may be used. Where the free acid is used, the reaction  
5 is preferably carried out in the presence of a condensing agent, for example: a disubstituted carbodiimide, such as N,N'-dicyclohexylcarbodiimide; an azolide  
40 compound, such as N,N'-carbonylimidazole; a dehydrating agent, such as N-ethoxycarbonyl-2-ethoxy-1,2-dihydro-quinoline, phosphorus oxychloride or an alkoxyacetylene,  
10 or a Vilsmeier reagent prepared from dimethylformamide and phosphorus oxychloride.

Where a reactive derivative of the acid of formula (VII) is employed, no such condensing agent is  
15 needed, but, depending upon the nature of the reactive derivative, the reaction may preferably be carried out in the presence of a base. Suitable bases include, for example: alkali metal compounds, such as sodium bicarbonate, potassium bicarbonate, sodium carbonate  
20 or potassium carbonate; and aliphatic, aromatic or nitrogen-containing heterocyclic bases, such as triethylamine, N,N-dimethylaniline, N,N-diethylaniline, N-methylpiperidine, N-methylpyrrolidine, pyridine, collidine or lutidine.

Preferred reactive derivatives of the acid (VII) include the acid halide, the acid anhydride, mixed acid anhydrides, active esters, active amides and the acid azide. Suitable mixed acid anhydrides include  
5 those with monoesters of carbonic acid (for example monomethyl carbonate or monoisobutyl carbonate) and those with lower alkanolic acids or lower haloalkanoic acids (such as pivalic acid or trichloroacetic acid). Suitable active esters include, for example, the p-  
10 nitrophenyl ester, the pentachlorophenyl ester, the N-hydroxyphthalimide ester and the N-hydroxybenzotriazole ester.

The reaction is preferably effected in the presence of a solvent, the nature of which is not  
15 critical, provided that it has no adverse effect upon the reaction. Suitable solvents include acetone, tetrahydrofuran, dioxan, ethyl acetate, chloroform, methylene chloride, dimethylformamide, acetonitrile and water, as well as mixtures of two or more of these  
20 solvents.

The reaction temperature is not particularly critical and the reaction is therefore normally performed at room temperature or with cooling. The time required for the reaction will vary, depending mainly on the

nature of the acylating agent and on the reaction temperature, but the reaction will normally be complete within from 10 minutes to several tens of hours.

5           Upon completion of the reaction, the desired compound of formula (VI) may be recovered from the reaction mixture by conventional means and, although the compound may, if necessary, be purified (for example by recrystallization or by the various chromatographic techniques) it may also be used, without  
10 ~~intermediate purification or separation~~, for the next step, that is to say the preparation of the desired compound of formula (I).

          The reaction to produce the compound of  
15 formula (I) comprises contacting the compound of formula (VI) with thiourea, preferably in the presence of a suitable solvent. The nature of the solvent is not critical, provided that it has no adverse effect upon the reaction. Suitable solvents include water,  
20 methanol, ethanol, dimethylformamide, dimethylacetamide, acetonitrile, tetrahydrofuran and mixtures of two or more of these solvents.

          If desired, a base (such as sodium acetate or sodium bicarbonate) may be added to the reaction mixture

in order to promote the reaction or assist it to go to completion. Formation of by-products may be prevented by effecting the reaction in the presence of a buffer solution of pH 6.5 - 7.  
Λ


5           The amount of thiourea employed is preferably 1 or more equivalents per equivalent of said compound of formula (VI).

          The reaction temperature is not particularly critical and the reaction is therefore preferably effected  
10 at ambient temperature. The time required for the reaction will vary, depending upon the reaction conditions, but a period of from several tens of minutes to several hours will generally be required.

          Upon completion of the reaction, the desired  
15 compound of formula (I) may be recovered by conventional means, for example by concentration under reduced pressure, extraction, reprecipitation or chromatography.


CL   Method 4

          P   Compounds of formula (I) may also be obtained  
20 by reacting a compound of formula (VIII):




15 (in which  $R^1$ ,  $R^2$  and  $R^3$  are as defined above) with thiourea.

1 Compounds of formula (VIII), which are new and also form part of the present invention, may be prepared by nitrosoating a compound of formula (IX):



14 (in which  $R^2$  and  $R^3$  are as defined above) to give a compound of formula (X):





(in which  $R^2$  and  $R^3$  are as defined above) and then alkylating the hydroxy group attached to the imino nitrogen atom of said compound of formula (X).

5 Representative examples of the new compounds  
of formula (VIII) include:

52. Acetoxymethyl 7-(4-chloro-2-methoxyimino-3-oxobutylaminol)-3-methoxymethyl-3-cephem-4-carboxylate

53. 1-Acetoxyethyl 7-(4-chloro-2-methoxyimino-3-oxobutylaminol-3-methoxymethyl-3-cephem-4-carboxylate

54. 1-Propionyloxymethyl 7-(4-chloro-2-methoxyimino-3-oxobutyrilaminol-3-methoxymethyl-3-cephem-4-carboxylate

15 55. 1-Ethoxycarbonyloxyethyl 7-(4-chloro-2-methoxy-  
imino-3-oxobutrylamino)-3-methoxymethyl-3-cephem-4-  
carboxylate

- P<sub>1</sub> 56. 1-Ethoxycarbonyloxyethyl 7-(4-chloro-2-ethoxy-  
imino-3-oxobutyrylamino)-3-methoxymethyl-3-cephem-4-  
carboxylate
- P<sub>1</sub> 57. Methoxycarbonyloxymethyl 7-(4-chloro-2-methoxy-  
5 imino-3-oxobutyrylamino)-3-methoxymethyl-3-cephem-4-  
carboxylate
- P<sub>1</sub> 58. Ethoxycarbonyloxymethyl 7-(4-chloro-2-ethoxy-  
imino-3-oxobutyrylamino)-3-methoxymethyl-3-cephem-4-  
carboxylate
- 10 P<sub>1</sub> 59. Isopropoxycarbonyloxymethyl 7-(4-chloro-2-  
methoxyimino-3-oxobutyrylamino)-3-methoxymethyl-3-  
cephem-4-carboxylate
- P<sub>1</sub> 60. Butoxycarbonyloxymethyl 7-(4-chloro-2-methoxy-  
imino-3-oxobutyrylamino)-3-methoxymethyl-3-cephem-4-  
15 carboxylate
- P<sub>1</sub> 61. 1-Propionyloxyethyl 7-(4-chloro-2-methoxy-  
imino-3-oxobutyrylamino)-3-methoxymethyl-3-cephem-4-  
carboxylate
- P<sub>1</sub> 62. 1-Butyryloxyethyl 7-(4-chloro-2-ethoxyimino-3-  
20 oxobutyrylamino)-3-methoxymethyl-3-cephem-4-carboxylate

P<sub>1</sub>

63. Isovaleryloxymethyl 7-(4-chloro-2-methoxy-  
imino-3-oxobutyrylamino)-3-methoxymethyl-3-cephem-4-  
carboxylate

P<sub>1</sub>

5

64. Pivaloyloxymethyl 7-(4-chloro-2-methoxyimino-  
3-oxobutyrylamino)-3-methoxymethyl-3-cephem-4-  
carboxylate

P<sub>1</sub>

65. Pivaloyloxymethyl 7-(4-chloro-2-ethoxyimino-  
3-oxobutyrylamino)-3-methoxymethyl-3-cephem-4-  
carboxylate

10 P<sub>1</sub>

66. Isobutyryloxymethyl 7-(4-chloro-2-methoxy-  
imino-3-oxobutyrylamino)-3-methoxymethyl-3-cephem-  
4-carboxylate

P<sub>1</sub>

15

67. 1-Pivaloyloxyethyl 7-(4-chloro-2-methoxy-  
imino-3-oxobutyrylamino)-3-methoxymethyl-3-cephem-4-  
carboxylate

P<sub>1</sub>

68. 1-(1-Ethylpropoxycarbonyloxy)ethyl 7-(4-chloro-  
2-methoxyimino-3-oxobutyrylamino)-3-methoxymethyl-3-  
cephem-4-carboxylate


P<sub>1</sub>

20

69. 3,3,3-Trimethylpropionyloxymethyl 7-(4-chloro-  
2-methoxyimino-3-oxobutyrylamino)-3-methoxymethyl-3-  
cephem-4-carboxylate.

37

The compound of formula (IX) can be prepared by acylating a compound of formula (III):



(in which  $R^2$  and  $R^3$  are as defined above) with 4-chloro-3-oxobutyl chloride (which can be obtained by reacting diketene with chlorinol). This acylation may be conducted by conventional means and is preferably effected in a solvent, the nature of which is not critical, provided that it has no adverse effect upon the reaction. Suitable solvents include methylene chloride, chloroform, tetrahydrofuran and dioxan. The acylation is preferably conducted in the presence of a base, preferably an organic base such as triethylamine, pyridine, N,N-dimethylaniline or N,N-diethylaniline. The reaction is preferably effected at about ambient temperature or at a lower temperature and will normally require a period of from several minutes to several hours. After completion of the reaction, the product of formula (IX) may be recovered and purified by

conventional means, for example concentration, extraction with organic solvents, chromatographic techniques or recrystallization.

*P* The nitrosoation of the compound of formula (IX) to prepare the compound of formula (X) may be effected by techniques known for the nitrosoation of reactive methylene groups. Such a nitrosoation reaction is normally effected using a metal salt of nitrous acid under acidic conditions or an ester of nitrous acid under suitable conditions. However, when preparing the compounds of the invention, it is necessary to carry out the reaction under such conditions that the cephalosporin ring system and the chlorine atom on the side chain at the 7- position do not participate in the reaction. It is, accordingly, desirable to carry out the reaction under weakly acidic or weakly basic conditions at a temperature below ambient. This reaction is normally carried out in the presence of a solvent, the nature of which is not critical, provided that it is capable of dissolving the compound of formula (IX) and does not have any adverse effect upon the reaction. Suitable solvents include formic acid, acetic acid, tetrahydrofuran, methanol, ethanol, chloroform, ethyl acetate and benzene, or a mixture of water with one or more of these solvents. The particular solvent chosen will depend upon the nature of the nitrosoating agent.

Examples of metal salts of nitrous acid employed as the nitrosoating agent include alkaline metal salts (such as sodium nitrite or potassium nitrite), preferably sodium nitrite. The nitrous acid ester is preferably an ester with a lower alcohol, for example pentyl nitrite or butyl nitrite.

Where a metal salt of nitrous acid is used as the nitrosoating agent, the reaction must be carried out under acidic conditions and, if an acidic solvent (such as formic acid or acetic acid) is not employed, the addition of an acid (which may be organic or inorganic) is necessary. Accordingly, we prefer to carry out the reaction using formic acid or acetic acid as the reaction solvent.

The reaction is preferably carried out at about ambient temperature or below and will require a period which may range from several minutes to several hours.

After completion of the reaction, the resulting product of formula (X) may be isolated and purified by conventional means, for example by concentration, extraction with organic solvents or chromatographic techniques.

The alkylation of the resulting compound of formula (X) to give the compound of formula (VIII) may

40

be effected by reacting the compound of formula (X) with an alkylating agent, preferably in the presence of a solvent. The nature of the solvent is not critical, provided that it has no adverse effect upon the reaction. Suitable solvents include acetone, tetrahydrofuran, dioxan, methanol, ethanol, chloroform, ethyl acetate, diethyl ether and dimethylformamide, or a mixture of two or more of these solvents.

Suitable alkylating agents include dialkyl sulphates (e.g. dimethyl sulphate or diethyl sulphate), diazoalkanes (e.g. diazomethane) and alkyl halides (e.g. methyl iodide or ethyl iodide).

Except when a diazoalkane (such as diazomethane) is used as the alkylating agent, the reaction is preferably effected in the presence of a base. Suitable bases include: alkali metal carbonates, such as sodium carbonate or potassium carbonate; alkali metal hydroxides, such as sodium hydroxide or potassium hydroxide; and nitrogen-containing organic bases, such as triethylamine, pyridine or N,N-dimethylaniline.

The reaction is preferably effected at ambient temperature or below and will normally require a period of from several minutes to several hours. After completion of the reaction, the desired compound of

formula (VIII) may be isolated and purified by conventional means, for example concentration, extraction with organic solvents, chromatographic techniques or recrystallization.

5           The reaction of the compound of formula (VIII) with thiourea to give the desired compound of formula (I) is essentially the synthesis of an aminothiazole derivative by reacting a haloketone with thiourea and may be carried out in much the same way as is common for this type of reaction.

10           The reaction is usually carried out in the presence of a solvent, the nature of which is not critical, provided that it has no adverse effect upon the reaction. The solvent is preferably an organic solvent (such as dimethylformamide, dimethylacetamide, methanol,  
15 ethanol or tetrahydrofuran) or a mixture of water with one or more of these organic solvents.

The thiourea is preferably employed in an amount of 1 or more equivalents per equivalent of said compound of formula (VIII).

20           In order to accelerate the reaction, sodium iodide may be added to the reaction mixture and the hydrogen chloride formed in the reaction may be neutralized by the addition of a neutral phosphate buffer solution.

42




The reaction is preferably effected at ambient temperature and will normally be complete within a period of from 1 to 10 hours.

When the reaction is complete, the desired compound of formula (I) may be isolated and purified by conventional means, for example by concentration, extraction with organic solvents, chromatographic techniques, reprecipitation or recrystallization.


CL Method 5

10 P Compounds of formula (I) may also be obtained by reacting a compound of formula (XI):



PS (in which  $R^2$ ,  $R^3$  and  $R^4$  are as defined above) with a compound of formula (XII): PS

15




TM (XII) PS

43

(in which  $R^1$  is as defined above) and then, if necessary, deprotecting the group represented by  $R^4$ .

Compounds of formula (XI) are new and also form part of the present invention. They may be prepared by reacting a compound of formula (XIII):



(XIII)

(in which  $R^4$  is as defined above) or a reactive derivative thereof with a compound of formula (III).

Representative examples of the novel compounds of formula (XI) include:

70. Pivaloyloxymethyl 7-[2-(2-aminothiazol-4-yl)-glyoxylamido]-3-methoxymethyl-3-cephem-4-carboxylate

71. Pivaloyloxymethyl 7-[2-(2-formamidothiazol-4-yl)glyoxylamido]-3-methoxymethyl-3-cephem-4-carboxylate

72. 1-Ethoxycarbonyloxyethyl 7-[2-(2-aminothiazol-4-yl)glyoxylamido]-3-methoxymethyl-3-cephem-4-carboxylate

73. 1-Ethoxycarbonyloxyethyl 7-[2-(2-formamido-  
thiazol-4-yl)glyoxylamido]-3-methoxymethyl-3-cephem-  
4-carboxylate.

In the reaction to produce the compound of  
formula (XI), the compound of formula (XIII) may be  
used either in the form of the free acid or in the form  
of a reactive derivative thereof. When the free acid  
is used, the reaction is preferably effected in the  
presence of a condensing agent, for example: a disub-  
stituted carbodiimide, such as N,N'-dicyclohexylcarbo-  
diimide; an imidazolidine, such as N,N'-carbonylimidazole  
or thionyl diimidazole; N-ethoxycarbonyl-2-ethoxy-1,2-  
dihydroquinoline; or a Vilsmeier reagent prepared from  
dimethylformamide and phosphorus oxychloride or thionyl  
chloride.

On the other hand, where a reactive derivative  
of the acid (XIII) is employed, there is no need to use  
a condensing agent, but, depending upon the nature of the  
reactive derivative, it may be preferred to effect the  
reaction in the presence of a base. Suitable bases  
include: alkali metal compounds, such as sodium  
bicarbonate, potassium bicarbonate, sodium carbonate or  
potassium carbonate; and aliphatic, aromatic or nitrogen-  
containing heterocyclic bases, such as triethylamine,  
N,N-dimethylaniline, N-methylpiperidine, N-methyl-

pyrrolidine, pyridine, collidine or lutidine.

Reactive derivatives of the acid (XIII) include the acid halides, the acid anhydride, mixed acid anhydrides, active esters, active amides and the acid azide. Examples of suitable mixed acid anhydrides include those with monoesters of carbonic acid (for example monomethyl carbonate or monoisobutyl carbonate) and those with lower alkanolic acids and lower halo-alkanoic acids (such as pivalic acid or trichloro-acetic acid). Suitable active esters include, for example, the p-nitrophenyl ester, the pentachloro-phenyl ester, the N-hydroxyphthalimide ester and the N-hydroxybenzotriazole ester.

The reaction is preferably effected in the presence of a solvent, the nature of which is not critical, provided that it has no adverse effect upon the reaction. Suitable solvents include acetone, methyl ethyl ketone, tetrahydrofuran, dioxan, ethyl acetate, chloroform, methylene chloride, dimethylformamide, acetonitrile and dimethyl sulphoxide, and mixtures of these solvents with water.

There is no particular limitation on the reaction temperature and accordingly the reaction is preferably effected at ambient temperature or with cooling.

4/11

The time required for the reaction will vary, depending mainly on the nature of the acylating method and on the reaction temperature, but it will normally require a period of from 10 minutes to several tens of hours.

5           After completion of the reaction, the compound of formula (XI) may be recovered from the reaction mixture by conventional means and it may, if desired then be purified by conventional techniques such as chromatography.

10           The reaction of the compounds of formulae (XI) and (XII) is normally performed in a solvent, the nature of which is not critical, provided that it has no adverse effect upon the reaction. Suitable solvents include dimethylformamide, dimethylacetamide, acetonitrile and  
15 various alcohols, as well as mixtures of these solvents with water.

The alkoxyamine of formula (XII) is preferably employed in the form of a salt with an inorganic acid (such as hydrochloric acid, nitric acid or sulphuric acid)  
20 or an organic acid (such as acetic acid or benzoic acid).

The reaction temperature is not critical, but we normally prefer to carry out the reaction at a temperature  
20 from ambient temperature to 60°C. The time required for

the reaction may vary, depending upon the reaction conditions, but will generally be from 10 minutes to several hours.

After completion of the reaction, the desired compound of formula (I) may be recovered from the reaction mixture by conventional means, for example by adding water and a water-immiscible solvent (such as ethyl acetate) to the reaction mixture, separating the organic layer under slightly alkaline conditions from the aqueous layer and then removing the organic solvent by distillation from this organic layer to give the desired compound.

Where the group  $R^4$  in the compound obtained by this process is a protected amino group, it may be deprotected using the techniques described in relation to Method 1.

The desired compound of formula (I) may, if necessary, be purified by conventional means such as recrystallization and/or chromatographic techniques.

The compounds of formula (I) and their acid addition salts may advantageously be employed in anti-bacterial compositions for oral administration. In order that a compound may be used for this purpose,

it is essential, as mentioned above, that it should be well absorbed through the digestive tract after oral administration. Good absorption through the digestive tract is demonstrated by a good recovery of the compound or of degradation products in the urine after oral  
5 administration.

The prior art compound (g) has a recovery rate in urine of 66.7%, which is very nearly comparable with the recovery rates of 75.9% and 78% of Compounds 5 and  
10 6, which are representative of the compounds of the present invention. These figures are quite satisfactory for the purposes of oral administration.

However, in addition to this good absorption through the digestive tract, it is desirable that  
15 compounds such as the prior art compound (g) and the compounds of the invention should, after hydrolyzation in vivo, be very active against gram-positive and gram-negative bacteria. The compounds of the invention, as well as compound (g), are hydrolyzed in vivo to  
20 the corresponding carboxylic acids and hence it is the antibacterial activities of these carboxylic acids, rather than of the esters, which are important from the clinical point of view. The activities of the carboxylic acids corresponding to Compounds No. 5 and 6 and to compound (g)  
25 against various bacteria are shown in the following

49

Table, in terms of their minimal inhibitory concentrations ( $\mu\text{g/ml}$ ).

Table

5

	Compound 5	Compound 6	Compound (g)
<u>Staphylococcus aureus</u> 209P	0.4	0.2	12.5
<u>Staphylococcus aureus</u> 56	0.8	0.4	25
<u>Escherichia coli</u> NIHJ	0.4	0.8	0.8
<u>Escherichia coli</u> 609	0.4	0.8	0.8
<u>Shigella flexneri</u> 2a	0.8	0.4	0.8
<u>Klebsiella pneumoniae</u> 806	0.1	0.2	0.2
<u>Klebsiella</u> sp. 846	0.8	0.8	1.5
<u>Proteus vulgaris</u>	0.01	0.01	<0.1
<u>Salmonella enteritidis</u> G.	0.2	0.4	0.4

10

15

20

70500x

50



a P It is clear from the above Table, that the <sup>compounds</sup> compounds of the invention and the prior art compound are all highly active against gram- negative bacteria, when administered orally.

However, whereas Compounds 5 and 6 are active against 5 Staphylococcus aureus, which is representative of the gram- positive bacteria, compound (g) has a rather low activity against these bacteria.

The compounds of the invention are preferably administered orally, for example in the form of capsules, 10 tablets, powders, syrups or suspensions. The dosage depends upon the age, symptoms and body weight of the patient and on the duration of treatment, but the dosage may normally range from 0.2 g to 5 g per day, preferably from 0.5 g to 3 g per day for adults; however, if 15 necessary, larger doses may be employed.

In the pharmaceutical compositions of the present invention, any conventional pharmaceutically acceptable carrier or diluent may be employed in admixture with the active compound or compounds. As the composition is 20 generally intended to be administered orally, it is desirably presented in a form readily absorbed through the stomach or intestines. Tablets or capsules are normally in unit dosage form and may contain binding agents (e.g. syrup, gum arabic, gelatin, sorbitol, gum tragacanth 25 or polyvinylpyrrolidone), diluents (e.g. lactose, sugar,

51

corn starch, calcium phosphate, sorbitol or glycine), lubricants, (e.g. magnesium stearate, talc, polyethylene glycol or silica), disintegrating agents (e.g. potato starch) or wetting agents (e.g. sodium lauryl sulphate) 5 or any combination thereof. The tablets may, if desired, be coated, e.g. with an enteric coating, as is well-known in the art.

Liquid formulations may be aqueous or oily suspensions, syrups, elixirs or similar compositions. 10 Alternatively, the composition may be a dried product which can then be redissolved in water or another suitable vehicle before administration. Such liquid formulations may contain conventional additives, such as suspending agents (e.g. sorbitol syrup, methylcellulose, 15 glucose/sugar syrup, gelatin, hydroxyethylcellulose, carboxymethylcellulose, aluminium stearate gel or hydrogenated edible fat), emulsifying agents (e.g. lecithin, monooleic acid sorbitol or gum arabic), non-aqueous vehicles (e.g. almond oil, fractionated coconut 20 oil, oily esters, propylene glycol or ethyl alcohol) or any combination of two or more thereof.

When the composition of the invention is formulated in unit dosage form, it preferably contains from 50 to 500 mg of the compound or compounds of the invention 25 per unit dose.

52

The preparation of the compounds of the present invention is further illustrated by the following Examples and the preparation of certain intermediates is illustrated by the following Preparations. The compounds of the invention are all in the syn configuration.

CC PREPARATION 1

CC Pivaloyloxymethyl 3-methoxymethyl-7-phenoxyacetamido-3-  
cephem-4-carboxylate

P 1 g of sodium 3-methoxymethyl-7-phenoxyacetamido-  
10 3-cephem-4-carboxylate was dissolved in 50 ml of dimethyl sulphoxide, and 975 mg of pivaloyloxymethyl bromide were added thereto, after which the mixture was stirred at room temperature for 15 minutes. The mixture was then diluted with 200 ml of ethyl acetate, washed in turn with  
15 50 ml of a saturated aqueous solution of sodium bicarbonate and 50 ml of a saturated aqueous solution of potassium bisulphate, and then dried over anhydrous magnesium sulphate. After filtering off the drying agent, the filtrate was concentrated under reduced pressure and the  
20 resulting residue was chromatographed through 100 g of silica gel eluted with a 1 : 1 by volume mixture of hexane and ethyl acetate, to afford 750 mg of the desired pivaloyloxymethyl 3-methoxymethyl-7-phenoxyacetamido-3-cephem-  
4-carboxylate.

Nuclear Magnetic Resonance spectrum ( $\text{CDCl}_3$ )  $\delta$  ppm:

- 1.25 (9H, singlet, t-butyl),  
 3.35 (3H, singlet,  $\text{OCH}_3$ ),  
 3.54 (2H, singlet, 2-cephem  $\text{H}_2$ ),  
 4.29 (2H, singlet,  $\text{CH}_2$  of methoxymethyl),  
 4.58 (2H, singlet,  $\text{CH}_2$  of phenoxyacetamido),  
 5.01 (1H, doublet,  $J = 5$  Hz, 6-cephem H),  
 5.6-6.1 (3H, multiplet, 7-cephem H and  $\text{CH}_2$  of pivaloyloxymethyl),  
 6.7-7.6 (6H, multiplet,  $\text{C}_6\text{H}_5$  and NH).

## PREPARATION 2

Pivaloyloxymethyl 7-amino-3-methoxymethyl-3-cephem-4-carboxylate p-toluenesulfonate

488 mg of phosphorus pentachloride were dissolved in 5 ml of dry methylene chloride, and then 120 mg of phosphorus oxychloride were added to the solution. Whilst the mixture was being stirred at room temperature, 247 mg of pyridine were added. The mixture was then cooled to  $-10^\circ\text{C}$ , and 769 mg of pivaloyloxymethyl 3-methoxymethyl-phenoxycetamido-3-cephem-4-carboxylate were added thereto. The temperature of the mixture was then allowed to rise gradually to room temperature. After stirring the mixture for 2 hours, it was again cooled to  $0^\circ\text{C}$ , and then 1.5 ml of propanol were added and the mixture again stirred for 30 minutes. A small amount of water was added to the

mixture, which was then stirred for a further 15 minutes. The mixture was diluted with 50 ml of ethyl acetate and washed with a saturated aqueous solution of sodium bicarbonate. The ethyl acetate layer was separated and dried over anhydrous magnesium sulphate. The drying agent was filtered off and the filtrate was concentrated by evaporation under reduced pressure. Diisopropyl ether was added to the residue and the wall of the vessel was scraped. The resulting precipitates were collected by filtration and dried to give 443 mg of the desired pivalo-yloxymethyl 7-amino-3-methoxymethyl-3-cephem-4-carboxylate. This compound was dissolved in 5 ml of ethyl acetate, and then an equimolar amount of p-toluenesulfonic acid monohydrate in 5 ml of ethyl acetate was added to the solution. The resulting mixture was allowed to stand at ambient temperature for 3 hours, affording 523 mg of the title compound melting at 160-170°C (with decomposition, recrystallized from methylene chloride and ethyl acetate) in the form of needles.

#### Elemental Analysis:

Calculated for  $C_{15}H_{22}N_2O_6S \cdot C_7H_8O_3S$ :

C, 49.80%; N, 5.70%; N, 5.28%; S, 12.08%.

Found: C, 49.76%; H, 5.60%; N, 5.00%; S, 12.06%.

#### PREPARATION 3

Benzhydryl 7-[2-(2-chloroacetamidothiazol-4-yl)-2-methoxyiminoacetamido]-3-methoxymethyl-3-cephem-4-carboxylate

To 0.057 ml of dimethylformamide were added 0.061 ml of phosphorus oxychloride, with ice-cooling and stirring. The mixture was then stirred at 40°C for 1 hour and then twice subjected to azeotropic distillation

with dry methylene chloride. 1 ml of ethyl acetate was added to the resulting mixture, which was then vigorously stirred at room temperature whilst 200 mg of 2-(2-chloroacetamidothiazol-4-yl)-2-methoxyiminoacetic acid were added. Stirring was continued for a further 30 minutes to give a mixture (a).

3120 10 Meanwhile, 200 mg of benzhydryl 7-amino-3-methoxymethyl-3-cephem-4-carboxylate and 145 mg of N,N-diethylaniline were dissolved in 5 ml of methylene chloride, and the mixture was stirred at -5°C to give a mixture (b).

Mixture (a) was then added dropwise to mixture (b) and the mixtures were stirred together for 15 minutes, after which the resulting reaction mixture was concentrated by evaporation under reduced pressure. 20 ml of ethyl acetate and 5 ml of water were then added to the residue and the ethyl acetate layer was separated. This layer was washed in turn with 5 ml of a saturated aqueous solution of sodium bicarbonate, 5 ml of a 5% w/v aqueous solution of hydrogen chloride and finally 5 ml of a saturated aqueous solution of sodium chloride, after which the solution was dried over anhydrous magnesium sulphate. The drying agent was filtered off and the filtrate was concentrated by evaporation under reduced pressure. The resulting residue was chromatographed

56

through 30 g of silica gel (Kieselgel 60), eluted with a 3 : 2 by volume mixture of hexane and ethyl acetate, to give 213 mg of the desired benzhydryl 7-[2-(2-chloroacetamidothiazol-4-yl)-2-methoxyiminoacetamido]-3-methoxymethyl-3-cephem-4-carboxylate.

Nuclear Magnetic Resonance spectrum ( $\text{CDCl}_3$ )  $\delta$  ppm:

3.19 (3H, singlet,  $\text{OCH}_3$  of methoxymethyl),  
 3.51 (2H, singlet, 2-cephem  $\text{H}_2$ ),  
 4.09 (3H, singlet,  $\text{OCH}_3$  of methoxyimino):  
 4.20 (2H, singlet,  $\text{CH}_2$  of methoxymethyl),  
 4.22 (2H, singlet,  $\text{CH}_2$  of chloroacetamido),  
 5.02 (1H, doublet,  $J = 5$  Hz, 6-cephem H),  
 5.86 (1H, doubled doublet,  $J = 5$  and 9 Hz, 7-cephem H),  
 6.7-7.6 (12H, multiplet).

#### ( ( PREPARATION 4

( ( (1) 7-[2-(2-Aminothiazol-4-yl)-2-methoxyiminoacetamido]-3-methoxymethyl-3-cephem-4-carboxylic acid trifluoroacetate

20 200 mg of benzhydryl 7-[2-(2-chloroacetamidothiazol-4-yl)-2-methoxyiminoacetamido]-3-methoxymethyl-3-cephem-4-carboxylate, followed by 45 mg of thiourea, were dissolved in 5 ml of dimethylacetamide. The solution was maintained at room temperature for 2 hours,

after which a saturated aqueous solution of sodium bicarbonate was added. The reaction mixture was then extracted with 20 ml of ethyl acetate and the extract was washed with water to remove excess thiourea and then dried over anhydrous magnesium sulphate. After the drying agent had been filtered off, the filtrate was concentrated by evaporation under reduced pressure. The resulting residue was chromatographed through 30 g of silica gel (Wacogel C-100), eluted with ethyl acetate, to afford 63 mg of benzhydryl 7-[2-(2-aminothiazol-4-yl)-2-methoxyiminoacetamido]-3-methoxymethyl-3-cephem-4-carboxylate.

The whole of this product was dissolved in 2 ml of anisole, and then 1 ml of trifluoroacetic acid was added to the solution, with ice-cooling and stirring. The mixture was then maintained at room temperature for 30 minutes, after which it was concentrated by evaporation under reduced pressure and diisopropyl ether was added to the residue. The resulting precipitates were collected by filtration and dried, to afford 27 mg of 7-[2-(2-aminothiazol-4-yl)-2-methoxyiminoacetamido]-3-methoxymethyl-3-cephem-4-carboxylic acid trifluoroacetate.

Nuclear Magnetic Resonance spectrum (deuteroacetone/D<sub>2</sub>O)

25  $\delta$  ppm: 3.29 (3H, singlet, OCH<sub>3</sub> of methoxymethyl);



57.

3.57 (2H, singlet, 2-cephem H<sub>2</sub>);  
3.96 (3H, singlet, OCH<sub>3</sub> of methoxyimino);  
4.27 (2H, singlet, CH<sub>2</sub> of methoxymethyl);  
5.15 (1H, doublet, J = 5.0 Hz, 6-cephem H);  
5 5.97 (1H, doublet, J = 5.0 Hz, 7-cephem H);  
6.59 (1H, singlet). <sup>32</sup> <sub>32</sub>

CL CL PREPARATION 5

CL 8 7-[2-(2-Chloroacetamidothiazol-4-yl)-2-methoxyimino-  
9 acetamido]-3-methoxymethyl-3-cephem-4-carboxylic acid

10 P A mixture of 7.65 g of benzhydryl 7-[2-(2-chloro-  
acetamidothiazol-4-yl)-2-methoxyiminoacetamido]-3-  
methoxymethyl-3-cephem-4-carboxylate, 25 ml of methylene  
chloride, 5 ml of anisole and 20 ml of trichloroacetic  
acid was allowed to react at room temperature for 30  
15 minutes. At the end of this time, 300 ml of diisopropyl  
ether were added to the reaction mixture and the resulting  
precipitates were collected by filtration, giving 5.95 g  
of 7-[2-(2-chloroacetamidothiazol-4-yl)-2-methoxyimino-  
acetamido]-3-methoxymethyl-3-cephem-4-carboxylic acid.

8 9  
20 Nuclear Magnetic Resonance spectrum (deuteroacetone/deutero-  
dimethyl sulphoxide)  $\delta$  ppm:

3.30 (3H, singlet, OCH<sub>3</sub> of methoxymethyl);  
3.60 (2H, singlet, 2-cephem H<sub>2</sub>);  
3.97 (3H, singlet, OCH<sub>3</sub> of methoxyimino);

59

- 4.25 (2H, singlet, CH<sub>2</sub> of methoxymethyl);  
 4.37 (2H, singlet, CH<sub>2</sub> of chloroacetamido);  
 5.20 (1H, doublet, 6-cephem H);  
 5.90 (1H, doubled doublet, J = 5.0 and 9.0 Hz,  
 5 7-cephem H);  
 7.40 (1H, singlet, 5-thiazole H);  
 9.50 (1H, doublet, J = 9 Hz, 7-cephem NH).

CL PREPARATION 6

CL89 7-[2-(2-Aminothiazol-4-yl)-2-ethoxyiminoacetamido]-3-  
 10 methoxymethyl-3-cephem-4-carboxylic acid trifluoro-  
acetate

Following the method of Preparation 3, 225 mg  
 of 2-(2-chloroacetamidothiazol-4-yl)-2-ethoxyimino-  
 acetic acid and 200 mg of benzhydryl 7-amino-3-methoxy-  
 15 methyl-3-cephem-4-carboxylate were reacted to give  
 280 mg of benzhydryl 7-[2-(2-chloroacetamidothiazol-4-yl)-  
 2-ethoxyiminoacetamido]-3-methoxymethyl-3-cephem-4-  
 carboxylate, in the form of a yellow powder.

Nuclear Magnetic Resonance spectrum (CDCl<sub>3</sub>) δ ppm:  
 1.28 (3H, triplet, OCH<sub>2</sub>CH<sub>3</sub>);  
 3.17 (3H, singlet, OCH<sub>3</sub>);  
 3.50 (2H, broad singlet, 2-cephem H<sub>2</sub>);  
 4.07 (2H, singlet, CH<sub>2</sub> of methoxymethyl);  
 4.0-4.5 (4H, multiplet, OCH<sub>2</sub>CH<sub>3</sub> and CH<sub>2</sub> of  
 chloroacetamido);

32 5.07 (1H, doublet,  $J = 5$  Hz, 6-cephem H);

5.93 (1H, doubled doublet,  $J = 5$  and 9 Hz,  
7-cephem H);

6.90 (1H, singlet, 5-thiazole H);

5 7.06 (1H, singlet, CH of benzhydryl);

89 7.31 (10H, singlet,  $(C_6H_5)_2$ );

32 8.10 (1H, doublet,  $J = 9$  Hz, 7-cephem NH).

191 mg of this benzhydryl 7-[2-(2-chloroacetamido-  
thiazol-4-yl)-2-ethoxyiminoacetamido]-3-methoxymethyl-  
10 3-cephem-4-carboxylate were then treated with 40 mg of  
thiourea, as described in Preparation 4, to give 117 mg  
of benzhydryl 7-[2-(2-aminothiazol-4-yl)-2-ethoxyimino-  
8 9 acetamido]-3-methoxymethyl-3-cephem-4-carboxylate, in  
the form of a pale pink powder, which was then treated  
15 with 1.5 ml of trifluoroacetic acid in a mixture of  
anisole and methylene chloride. When diisopropyl ether  
was added to the mixture, a precipitate was obtained and  
this was collected by filtration, to give 90 mg of  
7-[2-(2-aminothiazol-4-yl)-2-ethoxyiminoacetamido]-3-  
89 20 methoxymethyl-3-cephem-4-carboxylic acid trifluoro-  
acetate.

Nuclear Magnetic Resonance spectrum (deuterodimethyl  
sulphoxide)  $\delta$  ppm:

1.27 (3H, triplet,  $J = 7$  Hz,  $OCH_2CH_3$ );

3.23 (3H, singlet,  $OCH_3$ );

25

61

- 5 { 3.53 (2H, singlet, 2-cephem H<sub>2</sub>);  
 4.16 (2H, quartet, J = 7 Hz, OCH<sub>2</sub>CH<sub>3</sub>);  
 4.20 (2H, singlet, <sup>32</sup>CH<sub>2</sub> of methoxymethyl);  
 5.15 (1H, doublet, J = 5 Hz, 6-cephem H);  
 5.78 (1H, doubled doublet, <sup>32</sup>J = 5 and 9 Hz,  
 7-cephem H);  
 6.80 (1H, singlet, 5-thiazole H);  
 9.70 (1H, doublet, J = 9 Hz, 7-cephem NH);  
 8.5-10.0 (4H, broad multiplet, NH<sub>2</sub> and two COOH).

10

CL PREPARATION 7

CL t-Butyl 3-oxo-4-p-toluenesulphonyloxybutyrate

p To 50 ml of dry acetonitrile were added 7.1 g of t-butyl 4-bromo-3-oxobutyrate and 9.45 g of silver p<sup>⊖</sup> toluenesulphonate, and the mixture was stirred for 3 days  
 15 at room temperature, whilst shielding it from the light. The reaction mixture was then filtered and the filtrate was concentrated by evaporation in vacuo.

The resulting crystals containing an oily substance were dissolved in ethyl acetate and the insolubles were  
 20 removed by filtration. The filtrate was concentrated by evaporation in vacuo, to give a brown, oily substance, which was purified by column chromatography through silica gel, eluted with a 4 : 1 by volume mixture of cyclohexane and ethyl acetate. The resulting colourless, oily

62

substance was recrystallized from a 1 : 2 by volume mixture of diethyl ether and hexane, to afford 4.5 g of t-butyl 3-oxo-4-p-toluenesulphonyloxybutyrate, in the form of colourless prisms melting at 67 - 69°C.

5 Nuclear Magnetic Resonance spectrum (CDCl<sub>3</sub>) δ ppm:

1.43 (9H, singlet, t-butyl);

2.43 (3H, singlet, CH<sub>3</sub> of toluene);

3.43 (2H, singlet, -CH<sub>2</sub>COO-);

4.60 (2H, singlet, -SO<sub>2</sub>OCH<sub>2</sub>-);

7.20-7.90 (4H, C<sub>6</sub>H<sub>4</sub>).

Elemental Analysis:

Calculated for C<sub>15</sub>H<sub>20</sub>O<sub>6</sub>S:

C, 54.92%; H, 6.15%; S, 9.78%.

Found: C, 55.03%; H, 6.07%; S, 9.86%.

# PREPARATION 8

t-Butyl 2-hydroxyimino-3-oxo-4-p-toluenesulphonyloxy-  
butyrate

4.5 g of t-butyl 3-oxo-4-p-toluenesulphonyloxybutyrate were dissolved in 40 ml of acetic acid, and then 1.42 g of sodium nitrite were added, at room temperature, to the solution over a period of 10 minutes. The mixture was then stirred at room temperature for 50 minutes, after which 200 ml of ethyl acetate were added and the

63

mixture was then washed with an aqueous solution of sodium chloride. The ethyl acetate solution was dried over magnesium sulphate and, after filtering off the drying agent, the filtrate was concentrated by evaporation under reduced pressure to give a brown, oily substance. This oily substance was purified by column chromatography through silica gel, eluted with a 4 : 1 by volume mixture of cyclohexane and ethyl acetate, affording 1.66 g of t-butyl 2-hydroxyimino-3-oxo-4-p-toluene-sulphonyloxybutyrate, in the form of colourless crystals, melting at 106° - 108°C (with decomposition, recrystallized from a 1 : 1 by volume mixture of diethyl ether and petroleum ether).

15 { Nuclear Magnetic Resonance spectrum ( $\text{CDCl}_3$ )  $\delta$  ppm: 67  
 1.52 (9H, singlet, t-butyl);  
 2.43 (3H, singlet,  $\text{CH}_3$  of toluene);  
 5.04 (2H, singlet,  $-\text{SO}_2\text{OCH}_2\text{CO}-$ );  
 7.20-7.92 (4H,  $\text{C}_6\text{H}_4$ );  
 10.23 (1H, singlet, OH of hydroxyimino).

20 { Elemental Analysis:

Calculated for  $\text{C}_{15}\text{H}_{19}\text{NO}_7\text{S}$ :

C, 50.48%, H, 5.36%, N, 3.92%, S, 8.98%.

Found: C, 50.62%, H, 5.08%, N, 3.83%, S, 8.97%.

64

## CL PREPARATION 9

CL t-Butyl 2-methoxyimino-3-oxo-4-p-toluenesulphonyloxy-  
butyrate

P To an ice-cooled solution of 1.66 g of t-butyl  
5 2-hydroxyimino-3-oxo-4-p-toluenesulphonyloxybutyrate  
in 20 ml of dry acetone were added 960 mg of anhydrous  
potassium carbonate and 0.466 ml of dimethyl sulphate,  
and then the mixture was stirred at room temperature  
for 3 hours. The mixture was then poured into ice  
10 water and extracted with methylene chloride. The extract  
was washed with an aqueous solution of sodium chloride,  
dried over magnesium sulphate and concentrated by  
evaporation under reduced pressure to give a brown, oily  
substance. This was purified by column chromatography  
15 through silica gel, eluted with a 4 : 1 by volume mixture  
of cyclohexane and ethyl acetate, to afford 650 mg of  
t-butyl 2-(syn)-methoxyimino-3-oxo-4-p-toluenesulphonyl-  
oxybutyrate, as a pale yellow oil.

Nuclear Magnetic Resonance spectrum (CDCl<sub>3</sub>)  $\delta$  ppm:

20

1.50 (9H, singlet, t-butyl);

2.43 (3H, singlet, CH<sub>3</sub> of toluene);

4.07 (3H, singlet, OCH<sub>3</sub>);

5.05 (2H, singlet, -SO<sub>2</sub>OCH<sub>2</sub>CO-);

7.20-7.90 (4H, C<sub>6</sub>H<sub>4</sub>).

65

## CL PREPARATION 10

CL 2-Methoxyimino-3-oxo-4-p-toluenesulphonyloxybutyric acid

P To a solution of 478 mg of t-butyl 2-(syn)-methoxy-  
 5 imino-3-oxo-4-p-toluenesulphonyloxybutyrate in 1 ml of  
 methylene chloride were added 2 ml of trifluoroacetic  
 acid, and the mixture was stirred at room temperature  
 for 4 hours. The methylene chloride and the excess  
 trifluoroacetic acid were then distilled off in vacuo,  
 10 leaving a brown, oily substance, which was dissolved in  
 diisopropyl ether and allowed to stand, affording 178 mg  
 of 2-(syn)-methoxyimino-3-oxo-4-p-toluenesulphonyloxy-  
 butyric acid, in the form of colourless crystals melting  
 20 at 131 - 132°C (with decomposition).

## 15 Elemental Analysis:

P { Calculated for  $C_{12}H_{13}NO_7S$ :

C, 45.72%; H, 3.84%; N, 4.45%; S, 10.18%.

Found: C, 45.50%; H, 3.92%; N, 4.32%; S, 9.98%.

## 20 { Nuclear Magnetic Resonance spectrum (deuteroacetone)

$\delta$  ppm:

2.47 (3H, singlet,  $CH_3$  of toluene);

4.10 (3H, singlet,  $OCH_3$ );

5.20 (2H, singlet,  $-SO_2OCH_2CO-$ );

66



7.25-7.95 (4H, C<sub>6</sub>H<sub>4</sub>),

9.80 (1H, broad singlet, COOH).

# PREPARATION 11

5 Pivaloyloxymethyl 7-(2-methoxyimino-3-oxo-4-p-toluene-  
sulphonyloxybutyrylamino)-3-methoxymethyl-3-cephem-4-  
carboxylate

3120 To a suspension of 464 mg of 2-(syn)-methoxyimino-3-oxo-4-p-toluenesulphonyloxybutyric acid in 20 ml of methylene chloride, cooled to -5°C, was added 0.204 ml of triethylamine, and the mixture was stirred for 5 minutes, until completely dissolved. To the resulting solution were added 0.17 ml of oxalyl chloride and a drop of dimethylformamide and the mixture was stirred at -5°C for 20 minutes. On removing the solvent, there was left 2-(syn)-methoxyimino-3-oxo-4-p-toluenesulphonyloxy-15 butyryl chloride. This was dissolved in 10 ml of methylene chloride, and then 0.394 ml of N,N-diethylaniline, followed by the methylene chloride solution, were 3120 added, at -5°C, to a solution of 530 mg of pivaloyloxymethyl 7-amino-3-methoxymethyl-3-cephem-4-carboxylate p-toluenesulphonate in 20 ml of methylene chloride. This 20 mixture was stirred at -5°C for 5 minutes, after which 3120 the solvent was distilled off. The resulting residue was dissolved in ethyl acetate and washed with dilute aqueous hydrochloric acid. The ethyl acetate layer was

separated and dried over magnesium sulphate. After filtering off the drying agent, the filtrate was concentrated by evaporation under reduced pressure, to give a brown, oily substance. This was purified by column chromatography through silica gel eluted with a 4 : 1 by volume mixture of cyclohexane and ethyl acetate, to afford 510 mg of pivaloyloxymethyl 7-[2-(syn)-methoxyimino-3-oxo-4-p-toluenesulphonyloxybutyrylamino]-3-methoxymethyl-3-cephem-4-carboxylate, in the form of a colourless, foamy substance.

Nuclear Magnetic Resonance spectrum ( $\text{CDCl}_3$ )  $\delta$ , ppm:

1.22 (9H, singlet, t-butyl);

2.43 (3H, singlet,  $\text{CH}_3$  of toluene);

3.30 (3H, singlet,  $\text{OCH}_3$  of methoxymethyl);

3.51 (2H, singlet, 2-cephem  $\text{H}_2$ );

4.10 (3H, singlet,  $\text{OCH}_3$  of methoxyimino);

4.27 (2H, singlet,  $\text{CH}_2$  of methoxymethyl);

4.97 (1H, doublet,  $J = 5.0$  Hz, 6-cephem H);

5.07 (2H, singlet,  $-\text{SO}_2\text{OCH}_2\text{CO}-$ );

5.53-5.97 (3H, multiplet, 7-cephem H and  $-\text{OCH}_2-$  of pivaloyloxymethyl);

7.20-7.93 (5H, multiplet, 7-cephem NH and  $\text{C}_6\text{H}_4$ ).

68

CC PREPARATION 12

P Following the procedure described in Preparation 7, the following compounds were prepared:

P t-Butyl 4-methanesulphonyloxy-3-oxobutyrates,  
5 as a pale yellow oil.

10 P Nuclear Magnetic Resonance spectrum (CDCl<sub>3</sub>) δ ppm:  
 1.47 (9H, singlet, t-butyl); 67  
 3.14 (3H, singlet, CH<sub>3</sub>SO<sub>2</sub>);  
 3.45 (2H, singlet, -COCH<sub>2</sub>CO-);  
 4.87 (2H, singlet, -SO<sub>2</sub>OCH<sub>2</sub>CO-).  
 t-Butyl 4-ethanesulphonyloxy-3-oxobutyrates,  
 a yellow oil.

15 P Nuclear Magnetic Resonance spectrum (CDCl<sub>3</sub>) δ ppm:  
 1.32-1.62 (9H+3H, singlet + triplet, t-butyl +  
 N CH<sub>3</sub>CH<sub>2</sub>SO<sub>2</sub>); 30  
 3.30 (2H, quartet, J = 7.0 Hz, CH<sub>3</sub>CH<sub>2</sub>SO<sub>2</sub>);  
 3.47 (2H, singlet, -COCH<sub>2</sub>CO-); 32  
 4.87 (2H, singlet, -SO<sub>2</sub>OCH<sub>2</sub>CO-). 405

20 t-Butyl 4-benzenesulphonyloxy-3-oxobutyrates,  
 colourless needles melting at 94 - 96°C.  
 69

Nuclear Magnetic Resonance spectrum ( $\text{CDCl}_3$ )  $\delta$  ppm:

1.43 (9H, singlet, t-butyl);

3.43 (2H, singlet,  $-\text{COCH}_2\text{CO}-$ );

4.63 (2H, singlet,  $-\text{SO}_2\text{OCH}_2\text{CO}-$ );

7.40-8.03 (5H, multiplet,  $\text{C}_6\text{H}_5$ ).

### CL PREPARATION 13

Following the procedure described in Preparation 8, the following compounds were prepared:

t-Butyl 2-hydroxyimino-4-methanesulphonyloxy-

10 3-oxobutyrates, white crystals melting at  $103-104^\circ\text{C}$   
(with decomposition).

Nuclear Magnetic Resonance spectrum ( $\text{CDCl}_3$ /deutero-acetone)  $\delta$  ppm:

1.57 (9H, singlet, t-butyl);

3.20 (3H, singlet,  $\text{CH}_3$  of methanesulphonyl);

5.23 (2H, singlet,  $-\text{SO}_2\text{OCH}_2\text{CO}-$ );

11.93 (1H, singlet, OH of hydroxyimino).

t-Butyl 4-ethanesulphonyloxy-2-hydroxyimino-3-

oxobutyrates, colourless crystals melting at  $85-87^\circ\text{C}$

(with decomposition).

Nuclear Magnetic Resonance spectrum ( $\text{CDCl}_3$ )  $\delta$  ppm:

- 1.47 (3H, triplet,  $J = 7.0$  Hz,  $\text{CH}_3\text{CH}_2\text{SO}_2$ );  
 1.57 (9H, singlet, t-butyl), <sup>32</sup> *was*  
 3.33 (2H, quartet,  $J = 7.0$  Hz,  $\text{CH}_3\text{CH}_2\text{SO}_2$ );  
 5.23 (2H, singlet,  $-\text{SO}_2\text{OCH}_2\text{CO}-$ ), <sup>32</sup> *was*  
 5 10.50 (1H, singlet, OH of hydroxyimino).

t-Butyl 4-benzenesulphonyloxy-2-hydroxyimino-  
 3-oxobutyrate, colourless needles melting at  $93 - 95^\circ\text{C}$  <sup>20</sup>  
 (with decomposition).

*P* 10 } Nuclear Magnetic Resonance spectrum ( $\text{CDCl}_3$ )  $\delta$  ppm: <sup>67</sup>  
 1.57 (9H, singlet, t-butyl);  
 5.07 (2H, singlet,  $-\text{SO}_2\text{OCH}_2\text{CO}-$ );  
 7.40-8.03 (5H, multiplet,  $\text{C}_6\text{H}_5$ );  
 10.17 (1H, broad singlet, OH of hydroxyimino).

### *CL* PREPARATION 14

15 *P* Following the procedures described in Preparation 9, the following compounds were prepared:

*P* t-Butyl 4-methanesulphonyloxy-2-(syn)-methoxyimino-3-oxobutyrate, a colourless oil.

*P* 20 } Nuclear Magnetic Resonance spectrum ( $\text{CDCl}_3$ )  $\delta$  ppm: <sup>67</sup>  
 1.54 (9H, singlet, t-butyl);  
 3.19 (3H, singlet,  $\text{CH}_3$  of methanesulphonyl);

71

70.

4.10 (3H, singlet,  $\text{OCH}_3$  of methoxyimino);

5.23 (2H, singlet,  $-\text{SO}_2\text{OCH}_2\text{CO}-$ ).

t-Butyl 4-ethanesulphonyloxy-2-(syn)-methoxyimino-3-oxobutyrates, a pale yellow oil.

5 { Nuclear Magnetic Resonance spectrum ( $\text{CDCl}_3$ )  $\delta$  ppm:

1.43 (3H, triplet,  $J = 7.0$  Hz,  $\text{CH}_3\text{CH}_2\text{SO}_2$ );

1.50 (9H, singlet, t-butyl);

3.27 (2H, quartet,  $J = 7.0$  Hz,  $\text{CH}_2\text{CHSO}_2$ );

4.07 (3H, singlet,  $\text{OCH}_3$  of methoxyimino);

5.18 (2H, singlet,  $-\text{SO}_2\text{OCH}_2\text{CO}-$ ).

t-Butyl 4-benzenesulphonyloxy-2-(syn)-methoxyimino-3-oxobutyrates, a colourless oil.

15 { Nuclear Magnetic Resonance spectrum ( $\text{CDCl}_3$ )  $\delta$  ppm:

1.50 (9H, singlet, t-butyl);

4.05 (3H, singlet,  $\text{OCH}_3$  of methoxyimino);

5.07 (2H, singlet,  $-\text{SO}_2\text{OCH}_2\text{CO}-$ );

7.30-8.00 (5H, multiplet,  $\text{C}_6\text{H}_5$ ).

CC PREPARATION 15

Following the procedure described in Preparation 20. 10, the following compounds were prepared:

72

*P* 4-Methanesulphonyloxy-2-(syn)-methoxyimino-3-oxobutyric acid, a pale brown oil.

*P* Nuclear Magnetic Resonance spectrum (deuteroacetone)

$\delta$  ppm:

- 5* {
- 3.14 (3H, singlet, <sup>67</sup>CH<sub>3</sub> of methanesulphonyl);
  - 4.10 (3H, singlet, OCH<sub>3</sub> of methoxyimino);
  - 5.27 (2H, singlet, -SO<sub>2</sub>OCH<sub>2</sub>CO-);
  - 10.18 (1H, singlet, COOH).

4-Ethanesulphonyloxy-2-(syn)-methoxyimino-3-oxobutyric acid, melting at 85.5 - 89°C.

*P* Nuclear Magnetic Resonance spectrum (deuteroacetone)

$\delta$  ppm:

- 15* {
- 1.40 (3H, triplet, <sup>67</sup>J = 7.0 Hz, CH<sub>3</sub>CH<sub>2</sub>SO<sub>2</sub>);
  - 3.34 (2H, quartet, <sup>32</sup>J = 7.0 Hz, CH<sub>3</sub>CH<sub>2</sub>SO<sub>2</sub>);
  - 4.13 (3H, singlet, OCH<sub>3</sub> of methoxyimino);
  - 5.33 (2H, singlet, -SO<sub>2</sub>OCH<sub>2</sub>CO-);
  - 11.10 (1H, broad singlet, COOH).

4-Benzenesulphonyloxy-2-(syn)-methoxyimino-3-oxobutyric acid, as crystals.

*P* *20* Nuclear Magnetic Resonance spectrum (deuteroacetone)

$\delta$  ppm:

*67*  
*73*

- 4.06 (3H, singlet,  $\text{OCH}_3$  of methoxyimino);  
 5.17 (2H, singlet,  $-\text{SO}_2\text{OCH}_2\text{CO}-$ );  
 7.37-8.03 (5H, multiplet,  $\text{C}_6\text{H}_5$ );  
 10.33 (1H, singlet,  $\text{COOH}$ ).

5

# PREPARATION 16

Following the procedure described in Preparation 11, the following compounds were prepared:

Pivaloyloxymethyl 7-[4-methanesulphonyloxy-2-(syn)-methoxyimino-3-oxobutyrylamino]-3-methoxymethyl-

10. 3-cephem-4-carboxylate, a colourless, foamy substance.

Nuclear Magnetic Resonance spectrum ( $\text{CDCl}_3$ )  $\delta$  ppm:

- 1.21 (9H, singlet, t-butyl); 67  
 3.16 (3H, singlet,  $\text{CH}_3$  of methanesulphonyl);  
 3.30 (3H, singlet,  $\text{OCH}_3$  of methoxymethyl);  
 3.53 (2H, broad singlet, 2-cephem  $\text{H}_2$ );  
 4.13 (3H, singlet,  $\text{OCH}_3$  of methoxyimino);  
 4.24 (2H, singlet,  $\text{CH}_2$  of methoxymethyl);  
 4.99 (1H, doublet,  $J = 4.0$  Hz, 6-cephem H); 32  
 5.23 (2H, singlet,  $-\text{SO}_2\text{OCH}_2\text{CO}-$ );  
 5.60-5.93 (3H, multiplet, 7-cephem H and  $\text{CH}_2$  of pivaloyloxymethyl);  
 7.58 (1H, doublet,  $J = 9.0$  Hz, 7-cephem NH). 32



P  
Pivaloyloxymethyl 7-[4-ethanesulphonyloxy-2-  
(syn)-methoxyimino-3-oxobutyrylamino]-3-methoxymethyl  
3-cephem-4-carboxylate, a colourless, foamy substance.

Nuclear Magnetic Resonance spectrum (CDCl<sub>3</sub>) δ ppm:

- 5 {  
1.22 (9H, singlet, t-butyl); 67  
1.43 (3H, triplet, J = 7.0 Hz, CH<sub>3</sub>CH<sub>2</sub>SO<sub>2</sub>);  
3.27 (2H, quartet, J = 7.0 Hz, CH<sub>3</sub>CH<sub>2</sub>SO<sub>2</sub>);  
3.30 (3H, singlet, OCH<sub>3</sub> of methoxymethyl);  
3.54 (2H, broad singlet, 2-cephem H<sub>2</sub>);  
10 { 4.13 (3H, singlet, OCH<sub>3</sub> of methoxyimino);  
4.26 (2H, singlet, CH<sub>2</sub> of methoxymethyl);  
5.00 (1H, doublet, J = 5.0 Hz, 6-cephem H);  
5.27 (2H, singlet, -SO<sub>2</sub>OCH<sub>2</sub>CO-);  
15 { 5.60-5.97 (3H, multiplet, 7-cephem H and CH<sub>2</sub>  
of pivaloyloxymethyl);  
7.55 (1H, doublet, J = 9.0 Hz, 7-cephem NH).  
32

Pivaloyloxymethyl 7-[4-benzenesulphonyloxy-2-  
(syn)-methoxyimino-3-oxobutyrylamino]-3-methoxymethyl-  
3-cephem-4-carboxylate, a pale yellow, foamy substance.

20 { Nuclear Magnetic Resonance spectrum (CDCl<sub>3</sub>) δ ppm:

- 67  
1.22 (9H, singlet, t-butyl);  
3.30 (3H, singlet, OCH<sub>3</sub> of methoxymethyl);  
3.52 (2H, broad singlet, 2-cephem H<sub>2</sub>);  
75

- 4.10 (3H, singlet,  $\text{OCH}_3$  of methoxyimino);  
 4.27 (2H, singlet,  $\text{CH}_2$  of methoxymethyl);  
 4.98 (1H, doublet,  $J = 5.0$  Hz, 6-cephem H);  
 5.08 (2H, singlet,  $-\text{SO}_2\text{OCH}_2\text{CO}-$ );  
 5 5.60-5.90 (3H, multiplet, 7-cephem H and  $\text{CH}_2$   
 of pivaloyloxymethyl);  
 7.40-8.03 (6H, multiplet,  $\text{C}_6\text{H}_5$ , and 7-cephem NH).

CL PREPARATION 17

CL Pivaloyloxymethyl 7-(4-chloro-3-oxobutyrylamino)-3-  
 10 methoxymethyl-3-cephem-4-carboxylate

f 725 mg of diketene were dissolved in 10 ml of  
 dry methylene chloride and the solution stirred at  $-20^\circ\text{C}$ .  
 30 ml of a carbon tetrachloride solution containing 31 20  
 620 mg of chlorine were then added dropwise to the  
 15 solution, to produce 4-chloro-3-oxobutyryl chloride.

Meanwhile, 2 g of pivaloyloxymethyl 7-amino-3-  
 methoxymethyl-3-cephem-4-carboxylate p-toluenesulphonate  
 and 1.16 ml of N,N-diethylaniline were dissolved in  
 20 ml of methylene chloride. The resulting solution  
 20 was cooled to  $-10^\circ\text{C}$ , and then the 4-chloro-3-oxobutyryl  
 31 20  
 chloride solution obtained as described above was added  
 dropwise thereto. The mixture was then stirred at the  
 same temperature for 30 minutes, after which it was

26

concentrated by evaporation under reduced pressure. The resulting residue was dissolved in 50 ml of ethyl acetate and then washed in turn with water, a 5% w/v aqueous solution of hydrogen chloride and an aqueous solution of sodium chloride, after which it was dried over anhydrous magnesium sulphate and concentrated by evaporation under reduced pressure. The residue was dissolved in 3 ml of methylene chloride, and 30 ml of diethyl ether were added thereto, after which the mixture was allowed to stand. The resulting needle-like crystals were collected by filtration, washed with diethyl ether and dried to give 1.47 g of the title compound, melting at 131.5 - 132.5°C.

Nuclear Magnetic Resonance spectrum (CDCl<sub>3</sub>)  $\delta$  ppm:

- 1.23 (9H, singlet, t-butyl);  
 3.31 (3H, singlet, OCH<sub>3</sub>);  
 3.54 (2H, singlet, 2-cephem H<sub>2</sub>);  
 3.65 (2H, singlet, CH<sub>2</sub>);  
 4.26 (2H, singlet, CH<sub>2</sub>);  
 4.29 (2H, singlet, CH<sub>2</sub>);  
 4.97 (1H, doublet, J = 5.5 Hz, 6-cephem H);  
 5.65-6.0 (3H, multiplet, 7-cephem H and CH<sub>2</sub> of pivaloyloxymethyl);  
 7.64 (1H, doublet, J = 9 Hz, 7-cephem NH).

## CL PREPARATION 18

CL Pivaloyloxymethyl 7-(4-chloro-2-hydroxyimino-3-oxobutryl-  
amino)-3-methoxymethyl-3-cephem-4-carboxylate

P 2.57 g of pivaloyloxymethyl 7-(4-chloro-3-oxo-  
5 butrylamino)-3-methoxymethyl-3-cephem-4-carboxylate were  
dissolved in 25 ml of acetic acid, and then 409 mg of  
sodium nitrite were added, little by little, at room  
temperature to the solution, after which the mixture was  
stirred for 30 minutes. The mixture was then diluted  
10 with 200 ml of ethyl acetate, washed three times with a  
saturated aqueous solution of sodium chloride, dried over  
anhydrous magnesium sulphate and then concentrated by  
evaporation under reduced pressure. The residue was  
twice subjected to azeotropic distillation using toluene  
15 and the resulting residue was dried, giving 2.7 g of the  
title compound as a foamy solid.

Nuclear Magnetic Resonance spectrum (CDCl<sub>3</sub>)  $\delta$  ppm:

- 1.23 (9H, singlet, t-butyl), <sup>67</sup>  
3.33 (3H, singlet, OCH<sub>3</sub> of methoxymethyl);  
3.59 (2H, singlet, 2-cephem H<sub>2</sub>);  
4.33 (2H, singlet, CH<sub>2</sub> of methoxymethyl);  
4.75 (2H, singlet, ClCH<sub>2</sub>);  
5.05 (1H, doublet, J = 5.5 Hz, 6-cephem H);  
5.6-6.1 (3H, multiplet, <sup>32</sup>7-cephem H and CH<sub>2</sub> of  
pivaloyloxymethyl);  
9.3 (1H, doublet, J = 9 Hz, 7-cephem NH). <sup>32</sup>

## CL PREPARATION 19

CL Pivaloyloxymethyl 7-[4-chloro-2-(syn)-methoxyimino-3-oxobutyrylamino]-3-methoxymethyl-3-cephem-4-carboxylate

P 5 g of pivaloyloxymethyl 7-(4-chloro-2-hydroxy-  
 5 imino-3-oxobutyrylamino)-3-methoxymethyl-3-cephem-4-  
 carboxylate were dissolved in 40 ml of tetrahydrofuran.  
 To the resulting solution was added a solution of 2 g of  
 sodium carbonate in 40 ml of water, followed by 5 g of  
 dimethyl sulphate, after which the mixture was stirred  
 10 for 30 minutes. The mixture was then diluted with  
 150 ml of ethyl acetate, and washed twice with each in  
 turn of a saturated aqueous solution of sodium bicarbonate  
 and a saturated aqueous solution of potassium bisulphate,  
 after which it was dried over anhydrous magnesium  
 15 sulphate and concentrated by evaporation under reduced  
 pressure. The residue was chromatographed through 100 g  
 of silica gel eluted with a 3 : 1 by volume mixture of  
 chloroform and ethyl acetate, to give a solid containing  
 the title compound. This solid was dissolved in 30 ml  
 20 of diethyl ether and then left to stand under ice-  
 cooling, to produce crystals, which were washed with  
 diethyl ether and then dried, affording 1.9 g of the  
 title compound as needles melting at 168.5 - 169.5°C.

P Nuclear Magnetic Resonance spectrum (CDCl<sub>3</sub>) δ ppm:

25 1.24 (9H, singlet, t-butyl);

79

3.33 (3H, singlet, OCH<sub>3</sub> of methoxymethyl);  
 3.57 (2H, singlet, 2-cephem H<sub>2</sub>);  
 4.19 (3H, singlet, OCH<sub>3</sub> of methoxyimino);  
 4.30 (2H, singlet, CH<sub>2</sub> of methoxymethyl);  
 4.60 (2H, singlet, ClCH<sub>2</sub>);  
 5.03 (1H, doublet, J = 5.5 Hz, 6-cephem H);  
 5.6-6.1 (3H, multiplet, <sup>32</sup>7-cephem H and CH<sub>2</sub> of  
 N pivaloyloxymethyl);  
 7.19 (1H, doublet, J = 9 Hz NH).

10

C L PREPARATION 20

CL8 Pivaloyloxymethyl 7-[4-chloro-2-(syn)-ethoxyimino-3-  
 9 oxobutyrylamino]-3-methoxymethyl-3-cephem-4-carboxylate

p The procedure described in Preparation 19 was  
 repeated, but using diethyl sulphate in place of the  
 15 dimethyl sulphate. The title compound was obtained in  
 the form of needles melting at 171 - 172°C.

Nuclear Magnetic Resonance spectrum (CDCl<sub>3</sub>) δ ppm:

1.23 (9H, singlet, t-butyl); <sup>67</sup>  
 1.39 (3H, triplet, J = 7 Hz);  
 3.35 (3H, singlet, <sup>32</sup>OCH<sub>3</sub>);  
 3.57 (2H, singlet, 2-cephem H<sub>2</sub>);  
 4.32 (2H, singlet, CH<sub>2</sub> of methoxymethyl);  
 4.43 (2H, quartet, J = 7 Hz);  
 4.60 (2H, singlet, <sup>32</sup>ClCH<sub>2</sub>);  
 5.04 (1H, doublet, J = 5.5 Hz, 6-cephem H);  
<sup>32</sup>

20

25

80

+ 5.6-6.1 [3H, multiplet, 7-cephem H and CH<sub>2</sub> of pivaloyloxy-methyl]

7.17 (1H, doublet, J = 9 Hz, 7-cephem NH).

# PREPARATION 21

Pivaloyloxymethyl 7-[2-(2-formamidothiazol-4-yl)glyoxyl-  
amido]-3-methoxymethyl-3-cephem-4-carboxylate

To 0.544 ml of N,N-dimethylformamide was added, with ice-cooling, 0.582 ml of phosphorus oxychloride, and the resulting mixture was stirred at 40 - 45°C for 1 hour. The low boiling point materials were removed by allowing the mixture to stand for 5 minutes in vacuo, after which 10 ml of ethyl acetate, 1.25 g of 2-(2-formamidothiazol-4-yl)glyoxylic acid and 3 ml of N,N-dimethylformamide were added, in turn, to the resulting residue at room temperature. The mixture was stirred for 40 minutes and then added to a solution of 2.9 g of pivaloyloxymethyl 7-amino-3-methoxymethyl-3-cephem-4-carboxylate p-toluenesulphonate and 2.9 ml of N,N-diethylaniline in 30 ml of methylene chloride at a temperature of -20°C to -30°C. The mixture was then stirred at 0°C for 30 minutes, after which it was diluted with chloroform, washed in turn, with an aqueous solution of potassium bisulphite and an aqueous solution of sodium bicarbonate, and then dried over anhydrous magnesium sulphate. The solvent was removed by distillation and the residue was purified by column chromatography through

silica gel eluted with a 2 : 1 by volume mixture of ethyl acetate and chloroform, to give 1.9 g of the title compound in the form of an amorphous powder.

Nuclear Magnetic Resonance spectrum (deuterodimethylsulphoxide)  $\delta$  ppm:

- 5 { 1.22 (9H, singlet, t-butyl); 6.7  
3.32 (3H, singlet, OCH<sub>3</sub>);  
3.57 (2H, broad singlet, 2-cephem H<sub>2</sub>)  
4.32 (2H, broad singlet, CH<sub>2</sub> of methoxymethyl);  
5.07 (1H, singlet, 6-cephem H);  
10 { 5.7-6.0 (3H, multiplet, -COOCH<sub>2</sub>O<sup>2.1</sup> and 7-cephem H);  
8.03 (1H, broad doublet, <sup>3.2</sup>J = 9 Hz, 7-cephem NH);  
8.97 (1H, singlet);  
9.05 (1H, broad singlet).

#### PREPARATION 22

15 1-Ethoxycarbonyloxyethyl 7-[2-(2-formamidothiazol-4-yl)glyoxylamido]-  
3-methoxymethyl-3-cephem-4-carboxylate 9

The procedure described in Preparation 21 was repeated, but using 2.8 g of 1-ethoxycarbonyloxyethyl 7-amino-3-methoxymethyl-3-cephem-4-carboxylate p-toluenesulphonate and 1.25 g of 2-(2-formamido-20 thiazol-4-yl)glyoxylic acid, to give 1.5 g of the title compound.

Nuclear Magnetic Resonance spectrum (CDCl<sub>3</sub>)  $\delta$  ppm:

- 25 { 1.31 (3H, triplet, J = 7 Hz, OCH<sub>2</sub>CH<sub>3</sub>); 6.7  
1.59 (3H, doublet, J = 6 Hz, CH<sub>3</sub> of carbonyloxyethyl);  
3.32 (3H, singlet, OCH<sub>3</sub> of methoxymethyl);  
3.56 (2H, broad singlet, 2-cephem H);  
4.22 (1H, quartet; J = 7 Hz, OCH<sub>2</sub>CH<sub>3</sub>)  
4.32 (2H, singlet, CH<sub>2</sub> of methoxymethyl);  
5.03 (1H, doublet, J = 5 Hz, 6-cephem H);  
6.00 (1H, doubled doublet, J = 5+9 Hz, 7-cephem H);  
30 { 6.7-7.1 (1H, multiplet, CHCH<sub>3</sub>);  
7.38 (1H, singlet, 5-thiazole H);  
8.01 (1H, doublet, J = 9 Hz, 7-cephem H);  
8.60 (1H, singlet, HCO);  
9-12 (broad singlet (HCONH)).  
4.75

82



## DE CL EXAMPLE 1

CL8  
9 Pivaloyloxymethyl 7-[2-(2-aminothiazol-4-yl)-2-methoxy-  
iminoacetamido]-3-methoxymethyl-3-cephem-4-carboxylate

p To 71 mg of dimethylformamide were added, with  
5 ice-cooling and stirring, 135 mg of phosphorus oxychloride.  
20 The mixture was stirred at 40°C for 1 hour and then  
subjected twice to azeotropic distillation using dry  
methylene chloride. To the resulting mixture was added  
1 ml of ethyl acetate, after which, 265 mg of 2-(2-chloro-  
10 acetamidothiazol-4-yl)-2-methoxyiminoacetic acid were  
added, with vigorous stirring at room temperature, to the  
mixture and stirring was continued for 30 minutes.

Meanwhile, 121 mg of pivaloyloxymethyl 7-amino-  
3-methoxymethyl-3-cephem-4-carboxylate and 141 mg of  
15 N,N-diethylaniline were dissolved in 5 ml of methylene  
3120 chloride and stirred at -5°C. The resulting mixture was  
added dropwise to the mixture containing 2-(2-chloro-  
acetamidothiazol-4-yl)-2-methoxyiminoacetic acid prepared  
as described above. The reaction mixture was stirred  
20 for 15 minutes and then concentrated by evaporation under  
reduced pressure. To the residue were added 20 ml of  
ethyl acetate and 5 ml of water, and the ethyl acetate  
layer was separated, washed, in turn, with 5 ml of a  
saturated aqueous solution of sodium bicarbonate, 5 ml

of a 5% w/v aqueous solution of hydrogen chloride and 5 ml of a saturated aqueous solution of sodium chloride, and finally dried over anhydrous magnesium sulphate. The drying agent was filtered off and the filtrate was concentrated by evaporation under reduced pressure. The resulting residue was subjected to column chromatography through 10 g of silica gel eluted with a 2 : 1 by volume mixture of ethyl acetate and hexane, to afford 55 mg of pivaloyloxymethyl 7-[2-(2-chloroacetamido-  
8  
109 thiazol-4-yl)-2-methoxyiminoacetamido]-3-methoxymethyl-3-cephem-4-carboxylate.

This product was dissolved in 1 ml of dimethylacetamide, and 13.5 mg of thiourea were added to the resulting solution, which was then stirred at room  
15 temperature for 2 hours. The reaction mixture was then diluted with 20 ml of ethyl acetate, washed with a saturated aqueous solution of sodium bicarbonate and dried over anhydrous magnesium sulphate. The drying agent was filtered off and the filtrate was concentrated  
20 by evaporation under reduced pressure. The residue was subjected to column chromatography through 5 g of silica gel eluted with a 3 : 1 by volume mixture of ethyl acetate and hexane, to afford 36 mg of the title compound.

84

## Nuclear Magnetic Resonance spectrum (deuteroacetone)

$\delta$  ppm:  
 1.19 (9H, singlet, t-butyl),  
 3.23 (3H, singlet, OCH<sub>3</sub> of methoxymethyl),  
 3.52 (2H, singlet, 2-cephem H<sub>2</sub>),  
 3.90 (3H, singlet, OCH<sub>3</sub> of methoxyimino),  
 4.18 (2H, singlet, CH<sub>2</sub> of methoxymethyl),  
 5.12 (1H, doublet,  $J = 5$  Hz, 6-cephem H),  
 5.8-6.1 (3H, multiplet, 7-cephem H and CH<sub>2</sub> of  
 pivaloyloxymethyl),  
 6.78 (1H, singlet, 5-thiazole H),  
 6.6-7.1 (2H, broad singlet, NH<sub>2</sub>),  
 8.01 (1H, doublet,  $J = 9$  Hz, 7-cephem NH).

## EXAMPLE 2

Following the procedure described in Example 1, the following compounds were prepared:

Acetoxymethyl 7-[2-(2-aminothiazol-4-yl)-2-methoxyiminoacetamido]-3-methoxymethyl-3-cephem-4-carboxylate.

## Nuclear Magnetic Resonance spectrum (deuteroacetone)

$\delta$  ppm:  
 2.10 (3H, singlet, CH<sub>3</sub>CO),  
 3.22 (3H, singlet, OCH<sub>3</sub> of methoxymethyl),  
 3.52 (2H, singlet, 2-cephem H<sub>2</sub>),

3.92 (3H, singlet, OCH<sub>3</sub> of methoxyimino);  
 4.20 (2H, singlet, CH<sub>2</sub> of methoxymethyl);  
 5.11 (1H, doublet, J = 5 Hz, 6-cephem H);  
 5.6-6.3 (3H, multiplet, CH<sub>2</sub> of acetoxymethyl  
 and 7-cephem H);

6.76 (1H, singlet, 5-thiazole H);  
 6.6-7.1 (2H, broad singlet, NH<sub>2</sub>);  
 8.03 (1H, doublet, J = 9 Hz, 7-cephem NH).

Isovaleryloxymethyl 7-[2-(2-aminothiazol-4-yl)]

10 2-methoxyiminoacetamido]-3-methoxymethyl-3-cephem-4-carboxylate

Nuclear Magnetic Resonance spectrum (CDCl<sub>3</sub>) δ ppm:

0.99 (6H, doublet, J = 6.5 Hz, two CH<sub>3</sub> of  
 isovaleryl);  
 1.3-2.1 (1H, multiplet, CH of isovaleryl);  
 2.2-2.5 (2H, multiplet, CH<sub>2</sub> of isovaleryl);  
 3.32 (3H, singlet, OCH<sub>3</sub> of methoxymethyl);  
 3.56 (2H, broad singlet, 2-cephem H<sub>2</sub>);  
 3.98 (3H, singlet, OCH<sub>3</sub> of methoxyimino);  
 4.30 (2H, singlet, CH<sub>2</sub> of methoxymethyl);  
 5.06 (1H, doublet, J = 5.0 Hz, 6-cephem H);  
 5.8 (2H, broad singlet, NH<sub>2</sub>);  
 5.92 (2H, singlet, COOCH<sub>2</sub>OCO);  
 6.08 (1H, doubled doublet, J = 5.0 and 9.0 Hz,  
 7-cephem H);

86

6.70 (1H, singlet, 5-thiazole H);

8.20 (1H, doublet,  $J = 9.0$  Hz, 7-cephem NH).

32

Pivaloyloxymethyl 7-[2-(2-aminothiazol-4-yl)-  
2-ethoxyiminoacetamido]-3-methoxymethyl-3-cephem-  
5 carboxylate, having the properties described in Example 8.

### EXAMPLE 3

Following the procedure described in Example 1,  
1-ethoxycarbonyloxyethyl 7-[2-(2-aminothiazol-4-yl)-  
methoxyiminoacetamido]-3-methoxymethyl-3-cephem-  
10 carboxylate was prepared.

Nuclear Magnetic Resonance spectrum ( $\text{CDCl}_3$ )  $\delta$  ppm:

1.32 1.30 (3H, triplet,  $\text{OCH}_2\text{CH}_3$ );  
1.59 1.61 (3H, doublet,  $\text{CH}_3$  of carbonyloxyethyl);  
3.33 3.32 (3H, singlet,  $\text{OCH}_3$  of methoxymethyl);  
3.57 (2H, singlet, 2-cephem  $\text{H}_2$ );  
4.03 (3H, singlet,  $\text{OCH}_3$  of methoxyimino);  
4.23 4.21 (2H, quartet,  $\text{OCH}_2\text{CH}_3$ );  
4.34 4.30 (2H, singlet,  $\text{CH}_2$  of methoxymethyl);  
5.05 5.10 (1H, doublet,  $J = 5$  Hz, 6-cephem H);  
5.59 (1H, doubled doublet  $J = 5+9$  Hz, 7-cephem H);  
5.73 (2H, broad singlet  $\text{NH}_2$ );  
6.73 6.70 (1H, singlet, 5-thiazole H);  
6.7 - 7.1 (1H, multiplet, CH of ethoxycarbonyloxyethyl);  
7.90 (1H, doublet,  $J = 9$  Hz, 7-cephem NH).

CL EXAMPLE 4

CL Pivaloyloxymethyl 7-[2-(2-aminothiazol-4-yl)-2-methoxy-  
iminoacetamido]-3-methoxymethyl-3-cephem-4-carboxylate

8 9  
To 10 ml of dimethyl sulphoxide were added 1 g  
of 7-[2-(2-chloroacetamidothiazol-4-yl)-2-(syn)-methoxy-  
iminoacetamido]-3-methoxymethyl-3-cephem-4-carboxylic  
acid, 380 mg of bromomethyl pivalate and 240 mg of  
potassium fluoride, after which the mixture was stirred  
at room temperature for 1 hour. The mixture was then  
diluted with 100 ml of ethyl acetate and washed success-  
ively with water, a 5% w/v aqueous solution of sodium  
bicarbonate, a 10% w/v aqueous solution of potassium  
bisulphate and a saturated aqueous solution of sodium  
chloride, after which it was dried over anhydrous  
magnesium sulphate. The solvent was then distilled off  
under reduced pressure and the resulting residue was  
subjected to column chromatography through silica gel  
eluted with a 1 : 1 by volume mixture of chloroform and  
ethyl acetate, to give 300 mg of pivaloyloxymethyl 7-[2-  
(2-chloroacetamidothiazol-4-yl)-2-(syn)-methoxyimino-  
acetamido]-3-methoxymethyl-3-cephem-4-carboxylate as a  
pale yellow powder.

This compound was dissolved, with 60 mg of  
thiourea, in 3 ml of dimethylacetamide, and the solution

80

was stirred at room temperature for 4 hours. The mixture was then poured into 10 ml of a saturated aqueous solution of sodium bicarbonate and extracted with ethyl acetate. The extract was washed with, in turn, a 10% w/v aqueous solution of potassium bisulphate and a saturated aqueous solution of sodium chloride, after which it was dried over magnesium sulphate and concentrated by evaporation under reduced pressure. The residue was purified by column chromatography through silica gel eluted with a 3 : 1 by volume mixture of ethyl acetate and hexane to give 200 mg of the title compound. This compound was identified by nuclear magnetic resonance and found to be identical with the compound obtained in Example 1.

15

CL

EXAMPLE 5

Isobutyryloxymethyl 7-[2-(2-aminothiazol-4-yl)-2-methoxy-  
iminoacetamido]-3-methoxymethyl-3-cephem-4-carboxylate

p. 9

The procedure described in Example 4 was repeated, except that the bromomethyl pivalate was replaced by 360 mg of bromomethyl isobutyrate. There were obtained 180 mg of isobutyryloxymethyl 7-[2-(2-aminothiazol-4-yl)-2-(syn)-methoxyiminoacetamido]-3-methoxymethyl-3-cephem-4-carboxylate, as a slightly yellow powder.

89

88.

Nuclear Magnetic Resonance spectrum ( $\text{CDCl}_3$ )  $\delta$  ppm:

- 1.20 (6H, doublet,  $J = 6.5$  Hz, two  $\text{CH}_3$  of isobutyryl); <sup>32</sup>  
2.66 (1H, septet,  $J = 6.5$  Hz, CH of isobutyryl);  
3.21 (3H, singlet, <sup>32</sup> $\text{OCH}_3$  of methoxymethyl);  
3.40 (2H, AB quartet, 2-cephem  $\text{H}_2$ );  
4.01 (3H, singlet,  $\text{OCH}_3$  of methoxyimino);  
4.16 (2H, singlet,  $\text{CH}_2$  of methoxymethyl);  
5.05 (1H, doublet,  $J = 5$  Hz, 6-cephem H);  
5.6-6.2 (5H, multiplet, <sup>32</sup> $\text{NH}_2$ ,  $\text{CH}_2$  of carbonyloxy-  
N methyl and 7-cephem H);  
6.65 (1H, singlet, 5-thiazole H);  
8.06 (1H, doublet,  $J = 9$  Hz, 7-cephem NH). <sup>32</sup>

CL EXAMPLE 6

CL 15 8 Propionyloxymethyl 7-[2-(2-aminothiazol-4-yl)-2-methoxy-  
9 iminoacetamido]-3-methoxymethyl-3-cephem-4-carboxylate

P The procedure described in Example 4 was repeated except that the bromomethyl pivalate was replaced by 340 mg of bromomethyl propionate, to give 165 mg of the  
20 title compound as an almost colourless powder.

Nuclear Magnetic Resonance spectrum ( $\text{CDCl}_3$ )  $\delta$  ppm:

- 1.17 (3H, triplet,  $J = 6.5$  Hz,  $\text{CH}_2\text{CH}_3$ ); <sup>67</sup>  
2.41 (2H, quartet,  $J = 6.5$  Hz,  $\text{CH}_2\text{CH}_3$ ); <sup>32</sup> <sup>4.15</sup>

90



89.

3.20 (3H, singlet, CH<sub>3</sub> of methoxymethyl);  
3.35 (2H, AB quartet, 2-cephem H<sub>2</sub>);  
4.02 (3H, singlet, OCH<sub>3</sub> of methoxyimino);  
4.17 (2H, singlet, CH<sub>2</sub> of methoxymethyl);  
5.09 (1H, doublet, J<sub>32</sub> = 5 Hz, 6-cephem H);  
5.6-6.3 (5H, multiplet, NH<sub>2</sub>, CH<sub>2</sub> of carbonyloxy-  
N methyl and 7-cephem H);  
6.68 (1H, singlet, 5-thiazole H);  
8.25 (1H, doublet, J<sub>32</sub> = 9 Hz, 7-cephem NH).

CL EXAMPLE 7

CL<sub>8</sub> Pivaloyloxymethyl 7-[2-(2-aminothiazol-4-yl)-2-methoxy-  
9 iminoacetamido]-3-methoxymethyl-3-cephem-4-carboxylate

p To a solution of 45 mg of sodium 7-[2-(2-amino-  
thiazol-4-yl)-2-methoxyiminoacetamido]-3-methoxymethyl-  
15 3-cephem-4-carboxylate (prepared from the corresponding  
trifluoroacetate) in 1 ml of dimethylacetamide were

31 20 added, at -15°C, 27 mg of iodomethyl pivalate and the  
mixture was allowed to react for 15 minutes. At the  
end of this time, 20 ml of ethyl acetate were added to  
20 the reaction mixture, and the mixture was washed, in  
turn, with water, an aqueous solution of potassium  
bisulphate and an aqueous solution of sodium bicarbonate.  
The organic phase was separated and concentrated by  
evaporation under reduced pressure, and the residue was  
25 subjected to column chromatography through silica gel

91

eluted with a 3 : 1 by volume mixture of ethyl acetate and hexane, to give 49 mg of the title compound, whose properties were identical with those of the compound obtained in Example 1.

5

# CL EXAMPLE 8

Pivaloyloxymethyl 7-[2-(2-aminothiazol-4-yl)-2-ethoxy-  
iminoacetamido]-3-methoxymethyl-3-cephem-4-carboxylate

9

8  
109

The procedure described in Example 7 was repeated, except that sodium 7-[2-(2-aminothiazol-4-yl)-2-ethoxy-iminoacetamido]-3-methoxymethyl-3-cephem-4-carboxylate and iodomethyl pivalate were used, to give the title compound.

15

20

Nuclear Magnetic Resonance spectrum (CDCl<sub>3</sub>)  $\delta$  ppm:

- 1.22 (9H, singlet, t-butyl);
- 1.31 (3H, triplet, OCH<sub>2</sub>CH<sub>3</sub>);
- 3.30 (3H, singlet, OCH<sub>3</sub> of methoxymethyl);
- 3.53 (2H, singlet, 2-cephem H<sub>2</sub>);
- 4.28 (2H, quartet, OCH<sub>2</sub>CH<sub>3</sub>);
- 4.30 (2H, singlet, CH<sub>2</sub> of methoxymethyl);
- 5.01 (1H, doublet, J = 5 Hz, 6-cephem H);
- 5.7-6.2 (5H, multiplet, 7-cephem H, NH<sub>2</sub> and CH<sub>2</sub> of carbonyloxymethyl);
- 6.76 (1H, singlet, 5-thiazole H);
- 7.70 (1H, doublet, J = 9 Hz, 7-cephem NH).

92

## CL EXAMPLE 9

CL 8 1-Ethoxycarbonyloxyethyl 7-[2-(2-aminothiazol-4-yl)-2-ethoxyiminoacetamido]-3-methoxymethyl-3-cephem-4-carboxylate

- 5 P To a solution of 500 mg of sodium 7-[2-(2-aminothiazol-4-yl)-2-ethoxyiminoacetamido]-3-methoxymethyl-3-cephem-4-carboxylate in 5 ml of N,N-dimethylacetamide were added, with ice-cooling, 395 mg of 1-iodoethyl ethylcarbonate, and then the mixture was stirred at room
- 10 temperature for 30 minutes. At the end of this time, 50 ml of ethyl acetate were added to the reaction mixture, which was then washed with, in turn, 20 ml of water, 20 ml of a saturated aqueous solution of sodium bicarbonate and 20 ml of an aqueous solution of sodium chloride.
- 15 The mixture was then dried over anhydrous magnesium sulphate and the solvent was removed by distillation under reduced pressure, giving a residue, which was chromatographed through 20 g silica gel eluted with ethyl acetate, to afford 460 mg of the title compound.

20 Nuclear Magnetic Resonance spectrum (CDCl<sub>3</sub>)  $\delta$  ppm:

1.30 (3H, triplet, CH<sub>3</sub>CH<sub>2</sub>);

1.32 (3H, triplet, CH<sub>3</sub>CH<sub>2</sub>);

1.59 (3H, doublet, J = 6.0 Hz, CH<sub>3</sub> of carbonyloxyethyl;

25 3.30 (3H, singlet, OCH<sub>3</sub> of methoxymethyl);

- 3.52 (2H, broad singlet, 2-cephem H<sub>2</sub>);  
 4.22 (2H, quartet, CH<sub>3</sub>CH<sub>2</sub>);  
 4.27 (2H, quartet, CH<sub>3</sub>CH<sub>2</sub>);  
 4.30 (2H, singlet, CH<sub>2</sub> of methoxymethyl);  
 5 5.05 (1H, doublet, J = 5.0 Hz, 6-cephem H);  
 5.8 (2H, broad singlet, NH<sub>2</sub>);  
 6.00 (1H, doubled doublet, J 5.0 + 9.0 Hz,  
 7-cephem H);  
 6.75 (1H, singlet, 5-thiazole H);  
 10 6.7-7.1 (1H, multiplet, CH of carbonyloxyethyl);  
 7.8 (1H, doublet, J = 9 Hz, 7-cephem NH).

CL EXAMPLE 10

CL8  
9 Pivaloyloxymethyl 7-[2-(2-aminothiazol-4-yl)-2-methoxyiminoacetamido]-3-methoxymethyl-3-cephem-4-carboxylate

- 15 P To a solution of 510 mg of pivaloyloxymethyl  
 7-[2-(syn)-methoxyimino-3-oxo-4-p-toluenesulphonyloxy-  
 8 butyrylamino]-3-methoxymethyl-3-cephem-4-carboxylate in  
 9 5 ml of ethanol were added 76 mg of thiourea and 84 mg  
 of sodium acetate. 3 ml of water were then added drop-  
 20 wise to the mixture, after which the whole mixture was  
 stirred at room temperature for 3.5 hours. At the end  
 of this time, the ethanol was removed by distillation  
 and the residue was dissolved in ethyl acetate, washed  
 with an aqueous solution of sodium chloride and dried  
 25 over anhydrous magnesium sulphate. The ethyl acetate
- 94

was distilled off, giving a pale brown, foamy substance, which was purified by column chromatography through silica gel eluted with a 2 : 1 by volume mixture of ethyl acetate and methylene chloride, affording 392 mg of the title compound, in the form of a colourless foamy substance having the same properties as the product of Example 1.

CL EXAMPLE 11

CL 8 Propionyloxymethyl 7-[2-(2-aminothiazol-4-yl)-2-methoxy-  
9 iminoacetamido]-3-methoxymethyl-3-cephem-4-carboxylate.

10 P The procedure described in Example 10 was repeated,  
8 but using 490 mg of propionyloxymethyl 7-[2-(syn)-methoxy-  
9 imino-3-oxo-4-p-toluenesulphonyloxybutyrylamino]-3-methoxy-  
methyl-3-cephem-4-carboxylate, to give 370 mg of the  
title compound, having properties identical with those of  
15 the product of Example 6.

CL EXAMPLE 12

P The procedure described in Example 10 was repeated,  
8 except that the pivaloyloxymethyl 7-[2-(syn)-methoxyimino-  
9 3-oxo-4-p-toluenesulphonyloxybutyrylamino]-3-methoxymethyl-  
20 3-cephem-4-carboxylate was replaced by 1-ethoxycarbonyl-  
8 oxyethyl 7-[2-(syn)-methoxyimino-3-oxo-4-p-toluenesulphonyl-  
9 oxybutyrylamino]-3-methoxymethyl-3-cephem-4-carboxylate

95

8 or isobutyryloxymethyl 7-[2-(syn)-methoxyimino-3-oxo-4-  
 9 p-toluenesulphonyloxybutyrylamino]-3-methoxymethyl-3-  
 cephem-4-carboxylate, to give 1-ethoxycarbonyloxyethyl  
 8 9 7-[2-(2-aminothiazol-4-yl)-2-methoxyiminoacetamido]-3-  
 5 methoxymethyl-3-cephem-4-carboxylate (having properties  
 identical with those of the product of Example 3) and  
 8 isobutyryloxymethyl 7-[2-(2-aminothiazol-4-yl)-2-methoxy-  
 9 iminoacetamido]-3-methoxymethyl-3-cephem-4-carboxylate  
 (having properties identical with those of the product  
 10 of Example 5), respectively.

C L EXAMPLE 13

P The procedure described in Example 10 was  
 repeated, except that 465 mg of pivaloyloxymethyl   
 8 [4-methanesulphonyloxy-2-(syn)-methoxyimino-3-oxobutyryl-  
 15 9 amino]-3-methoxymethyl-3-cephem-4-carboxylate and 152 mg  
 of thiourea were used, to give 390 mg of pivaloyloxy-  
 8 9 methyl 7-[2-(2-aminothiazol-4-yl)-2-methoxyiminoacetamido]-  
 3-methoxymethyl-3-cephem-4-carboxylate, having properties  
 identical with those of the product of Example 1.

20 The same compound was also obtained following  
 the same procedure, but using, in separate experiments,  
 8 9 pivaloyloxymethyl 7-[4-ethanesulphonyloxy-2-(syn)-methoxy-  
 imino-3-oxobutyrylamino]-3-methoxymethyl-3-cephem-4-  
 9

- 8 carboxylate or pivaloyloxymethyl 7-[4-benzenesulphonyloxy-  
 9 2-(syn)-methoxyimino-3-oxobutyrylamino]-3-methoxymethyl-  
 3-cephem-4-carboxylate.

CL EXAMPLE 14

- CL 8 5 Pivaloyloxymethyl 7-[2-(2-aminothiazol-4-yl)-2-methoxy-  
 9 iminoacetamido]-3-methoxymethyl-3-cephem-4-carboxylate

P 47 mg of pivaloyloxymethyl 7-[4-chloro-2-(syn)-  
 methoxyimino-3-oxobutyrylamino]-3-methoxymethyl-3-  
 cephem-4-carboxylate were dissolved in 5 ml of dimethyl-  
 10 acetamide and then 14 mg of thiourea were added to the  
 solution, which was then stirred at room temperature for  
 4 hours. The reaction mixture was diluted with 50 ml  
 of ethyl acetate, washed three times, each time with  
 15 15 ml of water, dried over anhydrous magnesium sulphate  
 and then concentrated by evaporation under reduced pressure.  
 The resulting residue was dissolved in 1 ml of chloroform,  
 and 20 ml of diisopropyl ether were added to the  
 resulting solution. The precipitate produced was collected  
 by filtration and dried, to give 50 mg of the title  
 20 compound as a colourless powder having properties identical  
 with those of the product of Example 1.

## CL EXAMPLE 15

CL Pivaloyloxymethyl 7-[2-(2-aminothiazol-4-yl)-2-ethoxy-  
iminoacetamido]-3-methoxymethyl-3-cephem-4-carboxylate

P 9 The procedure described in Example 14 was

- 5 repeated, except that the pivaloyloxymethyl 7-[4-chloro-  
 2-(syn)-methoxyimino-3-oxobutyrylamino]-3-methoxymethyl-  
 3-cephem-4-carboxylate was replaced by<sup>9</sup> pivaloyloxy-  
 methyl 7-[4-chloro-2-(syn)-ethoxyimino-3-oxobutyrylamino]<sup>9</sup>  
 3-methoxymethyl-3-cephem-4-carboxylate, to give the  
 10 title compound as a colourless powder having properties  
 identical with those of the product of Example 8.

## CL EXAMPLE 16

CL Pivaloyloxymethyl 7-[2-(2-aminothiazol-4-yl)-2-methoxy-  
iminoacetamido]-3-methoxymethyl-3-cephem-4-carboxylate

- 15 P (a) A solution of 0.25 g of pivaloyloxymethyl 7-[2-(  
 (2-formamidothiazol-4-yl)glyoxylamido)-3-methoxymethyl-  
 3-cephem-4-carboxylate and 65 mg of methoxyamine hydro-  
 chloride in 2 ml of dimethylacetamide was stirred at  
 20 40°C for 140 minutes. At the end of this time, ethyl  
 20 acetate was added to the reaction mixture, which was  
 then washed with a saturated aqueous solution of sodium  
 chloride and dried over anhydrous magnesium sulphate.  
 The solvent was removed by distillation and the residue

98



was subjected to column chromatography through silica gel, eluted with a 2 : 1 by volume mixture of ethyl acetate and chloroform, to give 0.2 g of crude pivaloyloxymethyl 7-[2-(2-formamidothiazol-4-yl)-2-(syn)-methoxyiminoacetamido]-3-methoxymethyl-3-cephem-4-carboxylate, which was further purified by recrystallization from 1 ml of ethyl acetate, to give 170 mg of crystals melting at 172°C (with decomposition).

Nuclear Magnetic Resonance spectrum (deuterodimethyl sulphoxide)  $\delta$  ppm:

- 1.18 (9H, singlet, t-butyl), <sup>67</sup>
- 3.22 (3H, singlet, OCH<sub>3</sub> of methoxymethyl);
- 3.58 (2H, broad singlet, 2-cephem H<sub>2</sub>);
- 3.88 (3H, singlet, OCH<sub>3</sub> of methoxyimino);
- 4.14 (2H, singlet, CH<sub>2</sub> of methoxymethyl);
- 5.19 (1H, doublet, J = 5 Hz, 6-cephem H);
- 5.82 (3H, multiplet, <sup>32</sup>CH<sub>2</sub> of pivaloyloxymethyl and 7-cephem H);
- 7.37 (1H, singlet, 5-thiazole H);
- 8.47 (1H, singlet, HCO);
- 9.66 (1H, doublet, J = 9 Hz, 7-cephem NH);
- 12.58 (1H, broad singlet, <sup>32</sup>NH of formamido).

(b) To a solution of 2.6 g of the pivaloyloxymethyl 7-[2-(2-formamidothiazol-4-yl)-2-(syn)-methoxyiminoacetamido]-3-methoxymethyl-3-cephem-4-carboxylate prepared

999

as described above in 72 ml of methanol were added, with ice-cooling, 0.7 ml of concentrated hydrochloric acid, and the mixture was stirred at room temperature for 2.5 hours. The methanol was removed by distillation

5 in vacuo, and then 20 ml each of ethyl acetate and water were added to the residue, after which the mixture was neutralized by the addition of a saturated aqueous solution of sodium bicarbonate. The organic layer was washed with a saturated aqueous solution of sodium chloride, dried

10 and then concentrated by evaporation under reduced pressure. The residue was dissolved in 13 ml of chloroform and the solution was added dropwise, with stirring, to 100 ml of diisopropyl ether. The resulting precipitate was collected by filtration, to give 2.2 g of the title

15 compound in the form of a colourless powder whose properties were identical with those of the product of Example 1.

CL

EXAMPLE 17

CL

20 Pivaloyloxymethyl 7-[2-(2-aminothiazol-4-yl)-2-ethoxy-  
iminoacetamido]-3-methoxymethyl-3-cephem-4-carboxylate

p 9

The procedure described in Example 16(a) was repeated, except that the methoxyamine hydrochloride was replaced by 75 mg of ethoxyamine hydrochloride, to

8 give 150 mg of pivaloyloxymethyl 7-[2-(syn)-ethoxy-

(100)

9 imino-2-(2-formamidothiazol-4-yl)acetamido]-3-methoxymethyl-3-cephem-4-carboxylate, in the form of crystals  
 20 melting at 153°C.

Nuclear Magnetic Resonance spectrum (deuterodimethyl  
 (sulphoxide)  $\delta$  ppm:

5  
 1.18 (9H, singlet, t-butyl); 67  
 1.28 (3H, triplet,  $\text{OCH}_2\text{CH}_3$ );  
 3.21 (3H, singlet,  $\text{OCH}_3$  <sup>uns</sup> of methoxymethyl);  
 3.58 (2H, broad singlet, 2-cephem  $\text{H}_2$ );  
 10 4.15 (2H, singlet,  $\text{CH}_2$  of methoxymethyl);  
 4.19 (2H, quartet,  $\text{OCH}_2\text{CH}_3$ );  
 5.19 (1H, doublet,  $J = 5$  Hz, 6-cephem H);  
 5.71-5.95 (3H, multiplet,  $\text{CH}_2$  of pivaloyloxymethyl and 7-cephem H);  
 15 7.38 (1H, singlet, 5-thiazole H);  
 8.48 (1H, singlet, HCO);  
 9.64 (1H, doublet,  $J = 8$  Hz, 7-cephem NH);  
 12.60 (1H, broad singlet, NH of formamido);

The procedure described in Example 16(b) was  
 208 repeated, except that 9.65 g of pivaloyloxymethyl 7-[2-  
 9 (syn)-ethoxyimino-2-(2-formamidothiazol-4-yl)acetamido]-  
 3-methoxymethyl-3-cephem-4-carboxylate, 170 ml of  
 methanol and 2 ml of concentrated hydrochloric acid  
 were reacted at room temperature for 3 hours, to give

101

8.7 g of the title compound in the form of a colourless powder whose properties were identical to those of the product of Example 8.

CL EXAMPLE 18

CL 8  
9  
1-Ethoxycarbonyloxyethyl 7-[2-(2-aminothiazol-4-yl)-2-methoxyiminoacetamido]-3-methoxymethyl-3-cephem-4-carboxylate

8  
9  
10  
p A mixture of 180 mg of 1-ethoxycarbonyloxyethyl 7-[2-(2-formamidothiazol-4-yl)-2-(syn)-methoxyiminoacetamido]-3-methoxymethyl-3-cephem-4-carboxylate, 5 ml of methanol and 0.05 ml of concentrated hydrochloric acid were reacted as described in Example 16(b), to give 120 mg of the title compound, in the form of a pale yellow powder whose properties were identical with those of the product of Example 3.

15

CL EXAMPLE 19

CL  
Methoxycarbonyloxymethyl 7-[2-(2-aminothiazol-4-yl)-2-methoxyiminoacetamido]-3-methoxymethyl-3-cephem-4-carboxylate

20  
p To a solution of 500 mg of sodium 7-[2-(2-aminothiazol-4-yl)-2-(syn)-methoxyiminoacetamido]-3-methoxymethyl-3-cephem-4-carboxylate in 5 ml of dimethylacetamide

102

were added, with ice-cooling, 500 mg of iodomethyl methylcarbonate, and the mixture was stirred for 30 minutes. At the end of this time, the reaction mixture was diluted with 50 ml of ethyl acetate, washed, in turn, with a saturated aqueous solution of sodium bicarbonate and an aqueous solution of sodium chloride and dried over anhydrous magnesium sulphate. The magnesium sulphate was removed by filtration and the filtrate was concentrated by evaporation under reduced pressure. The residue was purified by column chromatography through silica gel, eluted with ethyl acetate, to give 433 mg of the title compound in the form of a foamy substance.

Nuclear Magnetic Resonance spectrum ( $\text{CDCl}_3$ )  $\delta$  ppm:

3.31 (3H, singlet,  $\text{OCH}_3$  of methoxymethyl);  
3.56 (2H, broad singlet, 2-cephem  $\text{H}_2$ );  
3.84 (3H, singlet,  $\text{OCH}_3$  of methoxycarbonyl);  
4.00 (3H, singlet,  $\text{OCH}_3$  of methoxyimino);  
4.31 (2H, singlet,  $\text{CH}_2$  of methoxymethyl);  
5.05 (1H, doublet, 6-cephem H);  
5.5-6.3 (5H, multiplet, 7-cephem H,  $\text{CH}_2$  of carbonyloxymethyl and  $\text{NH}_2$ );  
6.68 (1H, singlet, 5-thiazole H);  
8.10 (1H, doublet,  $J = 9.0$  Hz, 7-cephem NH).

32

103

## CL EXAMPLE 20

CL8  
9  
Ethoxycarbonyloxymethyl 7-[2-(2-aminothiazol-4-yl)-2-(syn)-methoxyiminoacetamido]-3-methoxymethyl-3-cephem-4-carboxylate

- 5 P To a solution of 861 mg of sodium 7-[2-(2-chloro-acetamidothiazol-4-yl)-2-(syn)-methoxyiminoacetamido]-3-methoxymethyl-3-cephem-4-carboxylate in 8.6 ml of 3120 dimethylacetamide were added, at -10°C, 565 mg of iodo-methyl ethylcarbonate and the mixture was stirred for 10 1 hour. At the end of this time, 100 ml of ethyl acetate were added to the reaction mixture, which was then washed, in turn, with water, a saturated aqueous solution of sodium bicarbonate and an aqueous solution of sodium chloride, and then dried over magnesium sulphate.
- 15 The organic layer was concentrated by evaporation under reduced pressure and the residue was purified by column chromatography through silica gel eluted with a 2 : 1 by volume mixture of ethyl acetate and chloroform, to 8 give 696 mg of ethoxycarbonyloxymethyl 7-[2-(2-chloro-acetamidothiazol-4-yl)-2-(syn)-methoxyiminoacetamido]-3-20 methoxymethyl-3-cephem-4-carboxylate.

The whole of this compound was dissolved in 6.4 ml of dimethylacetamide, and 800 mg of thiourea were added to the resulting solution, after which the mixture 25 was stirred at room temperature overnight. The mixture

104

was then diluted with 100 ml of ethyl acetate, washed three times with water and dried over anhydrous magnesium sulphate. The solvent was removed by distillation and the residue was subjected to column chromatography through silica gel eluted with ethyl acetate, to give 220 mg of the title compound in the form of a foamy substance.

Nuclear Magnetic Resonance spectrum ( $\text{CDCl}_3$ )  $\delta$  ppm:

- 1.32 (3H, triplet,  $J = 7$  Hz,  $\text{CH}_3$  of ethoxy);  
 3.32 (3H, singlet,  $\text{OCH}_3$  of methoxymethyl);  
 3.53 (2H, broad singlet, 2-cephem  $\text{H}_2$ );  
 3.98 (3H, singlet,  $\text{OCH}_3$  of methoxyimino);  
 4.23 (2H, quartet,  $J = 7$  Hz,  $\text{OCH}_2\text{CH}_3$ );  
 4.31 (2H, singlet,  $\text{CH}_2$  of methoxymethyl);  
 5.04 (1H, doublet,  $J = 6$  Hz, 6-cephem H);  
 5.6-6.3 (5H, multiplet, 7-cephem H,  $\text{CH}_2$  of carbonyloxymethyl and  $\text{NH}_2$ );  
 6.63 (1H, singlet, 5-thiazole H);  
 8.13 (1H, doublet,  $J = 9.0$  Hz, 7-cephem NH);

### EXAMPLE 21

Isovaleryloxymethyl 7-[2-(2-aminothiazol-4-yl)-2-methoxyiminoacetamido]-3-methoxymethyl-3-cephem-4-carboxylate

The procedure described in Example 20 was repeated to prepare the title compound, having the same properties as the second compound of Example 2.

C L EXAMPLE 22C L Capsules for oral administration

P The following mixture was compounded and  
encapsulated by conventional means with a No. 2 capsule,  
5 to give an encapsulated formulation:

	Pivaloyloxymethyl 7-[2-(2-amino- thiazol-4-yl)-2-( <u>syn</u> )-methoxyimino- acetamido]-3-methoxymethyl-3-cephem- 4-carboxylate	250 mg
10	Talc	5 mg
	Magnesium stearate	6.7 mg
	Sodium laurylsulphate	0.3 mg
	Lactose	28 mg

1060X

106